



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification n ⁶ : C07C 31/14, 31/16, C07D 213/24, A61K 31/095, 31/44		A1	(11) International Publication Number: WO 95/11883
			(43) International Publication Date: 4 May 1995 (04.05.95)

<p>(21) International Application Number: PCT/US94/08047</p> <p>(22) International Filing Date: 21 July 1994 (21.07.94)</p> <p>(30) Priority Data: 08/146,359 29 October 1993 (29.10.93) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 08/146,359 (CIP) Filed on 29 October 1993 (29.10.93)</p> <p>(71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): REITZ, David, B. [US/US]; 14814 Pleasant Ridge Court, Chesterfield, MO 63017 (US). LI, Jinglin [CN/US]; Apartment 306, 14550 Rialto Drive, Chesterfield, MO 63017 (US).</p> <p>(74) Agents: BULOCK, Joseph, W. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).</p>	<p>(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KE, KP, KR, KZ, LK, LT, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
--	--

(54) Title: **1,2-DIARYLCYCLOPENTENYL COMPOUNDS FOR THE TREATMENT OF INFLAMMATION**

(57) Abstract

A class of 1,2-diarylcyclopentenyl compounds is described for use in treating inflammation and inflammation-related disorders. Compounds of particular interest are defined by formula (III), wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkylcarbonyl and carboxyl; wherein R⁵ is selected from alkylsulfonyl, haloalkylsulfonyl and sulfamyl; and wherein B is phenyl or pyridyl, wherein B is optionally substituted at a substitutable position with alkyl, halo, alkylthio, cyano, haloalkyl, alkoxy, hydroxyalkyl and alkoxyalkyl; or a pharmaceutically-acceptable salt thereof.

(III)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

1,2 DIARYLCYCLOPENTENYL COMPOUNDS FOR THE TREATMENT OF INFLAMMATION

5

FIELD OF THE INVENTION

This invention is in the field of anti-inflammatory pharmaceutical agents and specifically relates to compounds, compositions and methods for treating inflammation and inflammation-associated disorders, such as arthritis.

BACKGROUND OF THE INVENTION

15

Prostaglandins play a major role in the inflammation process and the inhibition of prostaglandin production, especially production of PGG₂, PGH₂ and PGE₂, has been a common target of antiinflammatory drug discovery. However, common non-steroidal antiinflammatory drugs (NSAIDs) that are active in reducing the prostaglandin-induced pain and swelling associated with the inflammation process are also active in affecting other prostaglandin-regulated processes not associated with the inflammation process. Thus, use of high doses of most common NSAIDs can produce severe side effects, including life threatening ulcers, that limit their therapeutic potential. An alternative to NSAIDs is the use of corticosteroids, which have even more drastic side effects, especially when long term therapy is involved.

20

25

30

35

Previous NSAIDs have been found to prevent the production of prostaglandins by inhibiting enzymes in the human arachidonic acid/prostaglandin pathway, including the enzyme cyclooxygenase (COX). Recently, the sequence of another heretofore unknown enzyme in the human arachidonic acid/prostaglandin pathway has been reported by T. Hla and K. Nielson, *Proc. Natl. Acad. Sci, USA*, **89**, 7384 (1992) and named

"cyclooxygenase II (COX II)" or "prostaglandin G/H synthase II". The discovery of an inducible enzyme associated with inflammation provides a viable target of inhibition which more effectively reduces inflammation and produces fewer and less drastic side effects. Cyclooxygenase II is inducible by cytokines or endotoxins and such induction is inhibited by glucocorticoids (J. Masferrer, et al, *Proc. Natl. Acad. Sci, USA*, **89**, 3917 (1992)). The 6-methoxy-2-naphthylacetic acid metabolite of nabumetone has been found by E. Meade et al to selectively inhibit the COX II enzyme (*J. Biol. Chem.*, **268**, 6610 (1993)). In addition, Futaki et al (*Prostaglandins*, **47**, 55 (1994)) have reported that N-(2-cyclohexyloxy-4-nitrophenyl)methanesulfonamide inhibits the COX II enzyme.

The cyclopentene compounds disclosed herein are such safe and also effective antiinflammatory agents furthering such efforts. The invention compounds are found to show usefulness *in vivo* as antiinflammatory agents with minimal side effects. The substituted cyclopentene compounds disclosed herein preferably selectively inhibit cyclooxygenase II over cyclooxygenase I.

1,2-Diarylcycloalkenes have been made and used for a variety of utilities. For example, U.S. Patent No. 4,543,207 to Sato, et al., describes diphenyl cyclohexene as an electrical insulating oil additive.

U.S. Patent No. 4,229,207, to Laanio, et al., describes 1,2-diphenyl-cyclohex-1-ene-4-carboxylic acid esters, specifically, 1,2-di(4-phenylsulfonic acid)-cyclohex-1-ene-4-carboxylic acid ethyl ester. These ester compounds are reported to be effective as plant growth regulating agents and as post-emergent herbicides for controlling wild oats.

DE 4,212,628, published October 21, 1993, describes 3,4-bis-alkylphenylcyclohex-3-enes as anti-tumor agents.

5

The synthesis of 3,4-diphenyl- Δ^3 -cyclopentenone ethylene ketal is described as an intermediate for forming carbinols [E. J. Corey, et al., J. Amer. Chem. Soc., 85, 1788 (1963)]. 2,3-Bis-(4-hydroxyphenyl)-2-cyclopenten-1-one
10 has been identified from the knot resin powder of *Argaucaria angustifolia* [H. Ohash, et al., Phytochemistry, 31, 1371 (1992)].


Substituted 1,2-diphenylcyclopentenenes have been
15 synthesized for use in studies of their rotational behavior, includes specifically, 1-(2,4-Dimethylphenyl)-2-phenylcyclopentene [D. Y. Curtin, et al., J. Org. Chem., 36, 565 (1971)]. 1,2-Di-(2'-methoxyphenyl)- Δ^1 -cyclopentene has been identified as an impurity in the synthesis of
20 cannabinoids [O.P. Malik, et al., Ind. J. Chem., 14B, 975 (1976)].

1-(Substitutedphenyl)-2-phenylcyclopentenenes have been synthesized to study their photochemical reactions
25 into phenanthrene derivatives. Compounds with meta substituents, such as 1-(3-chlorophenyl)-2-phenylcyclopentene, are described in Somers, et al., J. Photochem. Photobiol., 48A, 353 (1989). Para substituents, including specifically 1-(4-fluorophenyl)-2-
30 phenylcyclopentene, are described in Laarhoven, Pure & Appl. Chem., 56, 1225 (1984).

DESCRIPTION OF THE INVENTION


35

A class of cyclopentene compounds useful in treating inflammation-related disorders is defined by Formula I:



I

wherein A is selected from




5

wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

wherein each of R³ through R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl;

provided that when A is



15

R⁵ cannot be hydrido when R¹⁰ is selected from hydrido, cyano, halo, methyl, trifluoromethyl and methoxy; further provided that R¹⁰ cannot be hydrido when R⁵ is selected from cyano, halo, methyl, trifluoromethyl and methoxy; and further provided that R⁵ and R¹⁰ are not both methoxy; or a pharmaceutically suitable salt thereof.

20

The phrase "further provided", as used in the above description, is intended to mean that the denoted proviso is not to be considered conjunctive with the other provisos.

25

Compounds of Formula I would be useful for, but not limited to, the treatment of inflammation in a subject, and for treatment of other inflammation-associated disorders, such as, as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. For example, compounds of Formula I would be useful to treat arthritis, including but not limited to rheumatoid arthritis, spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus and juvenile arthritis. Such compounds of Formula I would be useful in the treatment of asthma, bronchitis, menstrual cramps, tendinitis, bursitis, and skin related conditions such as psoriasis, eczema, burns and dermatitis. Compounds of Formula I also would be useful to treat gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome and ulcerative colitis and for the prevention of colorectal cancer. Compounds of Formula I would be useful in treating inflammation in such diseases as vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, scleroderma, rheumatic fever, type I diabetes, myasthenia gravis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, conjunctivitis, swelling occurring after injury, myocardial ischemia, and the like. The compounds are useful as anti-inflammatory agents, such as for the treatment of arthritis, with the additional benefit of having significantly less harmful side effects.

The present invention preferably includes compounds which selectively inhibit cyclooxygenase II over cyclooxygenase I and do not significantly inhibit one or more other arachidonic pathway steps, such as thromboxane B₂ (TXB₂) production.

Preferably, the compounds have a cyclooxygenase II IC₅₀ of less than about 0.1 μ M, and also have a selectivity ratio of cyclooxygenase II inhibition over cyclooxygenase I inhibition of at least 50, and more preferably of at least 100. Even more preferably, the compounds have a cyclooxygenase I IC₅₀ of greater than about 0.5 μ M, and more preferably of greater than 5 μ M. Such preferred selectivity may indicate an ability to reduce the incidence of common NSAID-induced side effects.


A preferred class of compounds consists of those compounds of Formula I wherein each of R¹ and R² is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxy carbonyl and carboxyl; and wherein each of R³ through R¹² is independently selected from hydrido, halo, lower alkyl, lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl.

A class of compounds of particular interest consists of those compounds of Formula I wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R³ through R¹² is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio,

cyano, fluoromethyl, difluoromethyl, trifluoromethyl,
 chloromethyl, dichloromethyl, trichloromethyl,
 pentafluoroethyl, heptafluoropropyl, difluorochloromethyl,
 dichlorofluoromethyl, difluoroethyl, difluoropropyl,
 5 dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy,
 butoxy, isopropoxy, tert-butoxy, methylenedioxy,
 trifluoromethoxy, hydroxymethyl, methoxymethyl,
 ethoxymethyl, isopropoxymethyl, tert-butoxymethyl,
 propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl,
 10 methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and
 sulfamyl.

Within Formula I there is a subclass of compounds
 of high interest represented by Formula II:

15



II

wherein each of R¹ and R² is independently
 selected from alkyl, hydrido, hydroxyalkyl, halo,
 20 haloalkyl, alkoxycarbonyl and carboxyl; and
 wherein each of R³ through R¹² is independently
 selected from hydrido, halo, alkyl, alkylthio, cyano,
 haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl,
 alkylsulfonyl, haloalkylsulfonyl and sulfamyl;
 25 provided that R⁵ cannot be hydrido when R¹⁰ is
 selected from hydrido, cyano, halo, methyl, trifluoromethyl
 and methoxy; further provided that R¹⁰ cannot be hydrido
 when R⁵ is selected from cyano, halo, methyl,
 trifluoromethyl and methoxy; and further provided that R⁵
 30 and R¹⁰ are not both methoxy;
 or a pharmaceutically suitable salt thereof.

A preferred class of compounds consists of those compounds of Formula II wherein each of R^1 and R^2 is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxy carbonyl and carboxyl; and wherein each of R^3 through R^{12} is independently selected from hydrido, halo, lower alkyl, lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl.

10

A class of compounds of particular interest consists of those compounds of Formula II wherein each of R^1 and R^2 is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R^3 through R^{12} is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl, methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl.

35

A more preferred class of compounds consists of those compounds of Formula II wherein R⁵ is selected from methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl.

5

A family of specific compounds of particular interest within Formula II consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 10 1-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
15 1-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
20 1-[2-(3,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
25 1-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
30 1-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
35 1-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-[3,4-di(difluoromethyl)phenyl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
5 4-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-benzenesulfonamide;
4-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
30 4-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
35 4-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
5 4-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-[3,4-di(difluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
30 4-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
35 4-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;

1-(2-phenylcyclopenten-1-yl)-4-(methylsulfonyl)benzene;

- 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(4-bromophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-iodophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-ethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-cyanophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2-methyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2-methyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(2-fluoro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(2-fluoro-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2-chloro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(2-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
5 4-(2-phenylcyclopenten-1-yl)benzenesulfonamide;
4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-cyanophenyl)cyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-methylthiophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(2-methyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-methyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(2-fluoro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-fluoro-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
30 4-[2-(2-chloro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
35 4-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;

- 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(fluoromethylsulfonyl)benzene;
- 1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 5-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
- 5-[2-(4-(aminosulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
- 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(4-trifluoromethylphenyl)-4,4-diethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene ;
- 35 1-[2-(4-trifluoromethylphenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide;
5 4-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(4-trifluoromethylphenyl)-4,4-dicarbomethoxycyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(4-trifluoromethylphenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
1-[2-(4-fluorophenyl)-4-methylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
30 1-[2-(4-fluorophenyl)-4-ethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4-(hydroxymethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
35 1-[2-(4-fluorophenyl)-4-carbomethoxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4-fluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(4-fluorophenyl)-4-(trifluoromethylcyclopenten-1-yl)]-4-(methylsulfonyl)benzene;
- 1-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(4-fluorophenyl)-4-(fluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4-methylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4-ethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(4-trifluoromethylphenyl)-4-(hydroxymethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4-carbomethoxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(4-trifluoromethylphenyl)-4-fluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4-(trifluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(4-trifluoromethylphenyl)-4-(fluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2,4,6-trifluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 4-[2-(4-fluorophenyl)-4-methylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4-ethylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4-(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
- 30 4-[2-(4-fluorophenyl)-4-carbomethoxycyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4-fluorocyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-(4-fluorophenyl)-4-(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(4-fluorophenyl)-4-(fluoromethyl)cyclopenten-1-yl]
benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4-methylcyclopenten-1-yl]
benzenesulfonamide;
5 4-[2-(4-trifluoromethylphenyl)-4-ethylcyclopenten-1-yl]
benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4-(hydroxymethyl)
cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4-carbomethoxycyclopenten-1-
10 yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4-fluorocyclopenten-1-yl]
benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4-
trifluoromethylcyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(2,4,6-trifluorophenyl)cyclopenten-1-
yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]
benzenesulfonamide; and
4-[2-(4-trifluoromethylphenyl)-4-(fluoromethyl)cyclopenten-
20 1-yl]benzenesulfonamide.


A family of compounds of more particular interest within Formula II consists of compounds and their pharmaceutically-acceptable salts as follows:

- 25 1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
30 1-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
4-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]
35 benzenesulfonamide;
1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
4-[2-(4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
- 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(fluoromethylsulfonyl)benzene;
- 5 5-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
- 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-(trifluoromethyl)phenyl)cyclopenten-1-yl]benzenesulfonamide;
- 1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 4-[2-(2,3-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 1-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 4-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 1-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 25 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 4-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 30 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 1-[2-(3,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 4-[2-(3,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;

- 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 5 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 10 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 1-(2-phenylcyclopenten-1-yl)-4-(methylsulfonyl)benzene;
 1-[2-(4-cyanophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 15 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 1-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
 20 1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene; and
 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene.

25 Within Formula I there is a second subclass of compounds of high interest represented by Formula III:



III

30 wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl;

wherein R⁵ is selected from alkylsulfonyl, haloalkylsulfonyl and sulfamyl; and

wherein B is phenyl or pyridinyl, wherein B is optionally substituted at a substitutable position with
5 alkyl, halo, alkylthio, cyano, haloalkyl, alkoxy, hydroxyalkyl and alkoxyalkyl; or a pharmaceutically-acceptable salt thereof.


A preferred class of compounds consists of those
10 compounds of Formula III wherein each of R¹ and R² is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxy carbonyl and carboxyl; wherein R⁵ is selected from lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl; and
15 wherein B is phenyl or pyridinyl, wherein B is optionally substituted at a substitutable position with lower alkyl, halo, lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower hydroxyalkyl and lower alkoxyalkyl.

20 A class of compounds of particular interest consists of those compounds of Formula III wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo,
25 iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl,
30 ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; wherein R⁵ is selected from methylsulfonyl, trifluoromethylsulfonyl and sulfamyl; and wherein B is optionally substituted with a radical selected
35 from fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl,

trifluoromethyl, chloromethyl, dichloromethyl,
 trichloromethyl, pentafluoroethyl, heptafluoropropyl,
 difluorochloromethyl, dichlorofluoromethyl, difluoroethyl,
 difluoropropyl, dichloroethyl, dichloropropyl, methoxy,
 5 ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy,
 methylenedioxy, trifluoromethoxy, hydroxymethyl,
 methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-
 butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl
 and pentoxymethyl.

10

Within Formula I there is a third subclass of
 compounds of high interest represented by Formula IV:



IV

15

wherein each of R¹ and R² is independently
 selected from alkyl, hydrido, hydroxyalkyl, halo,
 haloalkyl, alkoxycarbonyl and carboxyl; and

20 wherein each of R³ through R⁷ and R⁹ through R¹²
 is independently selected from hydrido, halo, alkyl,
 alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy,
 hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl
 and sulfamyl; or a pharmaceutically-acceptable salt
 thereof.

25

A preferred class of compounds consists of those
 compounds of Formula IV wherein each of R¹ and R² is
 independently selected from lower alkyl, hydrido, lower
 hydroxyalkyl, halo, lower haloalkyl, lower alkoxycarbonyl
 30 and carboxyl; and wherein each of R³ through R⁷ and R⁹
 through R¹² is independently selected from hydrido, halo,
 lower alkyl, lower alkylthio, cyano, lower haloalkyl, lower

alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl.

5 A class of compounds of particular interest consists of those compounds of Formula IV wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, 10 iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, 15 ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R³ through R⁷ and R⁹ through R¹² is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, 20 propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, 25 dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, 30 propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl, methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl.

35 A family of specific compounds of particular interest within Formula IV consists of compounds and pharmaceutically-acceptable salts thereof as follows:

2-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]pyridine;

- 5-fluoro-2-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]pyridine;
5-chloro-2-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]pyridine;
5 5-methyl-2-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]pyridine;
5-cyano-2-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]pyridine;
10 4-[2-(pyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-fluoropyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-chloropyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(5-methylpyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-cyanopyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(5-trifluoromethylpyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-methylcyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-ethylcyclopenten-1-yl]pyridine;
25 5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-(hydroxymethyl)cyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-carbomethoxycyclopenten-1-yl]pyridine;
30 5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-fluorocyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)cyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-carboxycyclopenten-1-yl]pyridine;
35 5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-(fluoromethyl)cyclopenten-1-yl]pyridine;

- 5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-methylcyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-ethylcyclopenten-1-yl]pyridine;
5 5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-(hydroxymethyl)cyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-carbomethoxycyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-fluorocyclopenten-1-yl]pyridine;
10 5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)cyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-carboxycyclopenten-1-yl]pyridine;
15 5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4-(fluoromethyl)cyclopenten-1-yl]pyridine;
4-[2-(5-fluoropyridin-2-yl)-4-methylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-fluoropyridin-2-yl)-4-ethylcyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(5-fluoropyridin-2-yl)-4-(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-fluoropyridin-2-yl)-4-carbomethoxycyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(5-fluoropyridin-2-yl)-4-fluorocyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-fluoropyridin-2-yl)-4-(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-fluoropyridin-2-yl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
30 4-[2-(5-fluoropyridin-2-yl)-4-(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-trifluoromethylpyridin-2-yl)-4-methylcyclopenten-1-yl]benzenesulfonamide;
35 4-[2-(5-trifluoromethylpyridin-2-yl)-4-ethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-trifluoromethylpyridin-2-yl)-4-(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(5-trifluoromethylpyridin-2-yl)-4-carbomethoxycyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-trifluoromethylpyridin-2-yl)-4-fluorocyclopenten-1-yl]benzenesulfonamide;
5 4-[2-(5-trifluoromethylpyridin-2-yl)-4-(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-trifluoromethylpyridin-2-yl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-trifluoromethylpyridin-2-yl)-4-fluoromethylcyclopenten-1-yl]benzenesulfonamide;
10 5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4,4-diethylcyclopenten-1-yl]pyridine;
15 5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4-di(hydroxymethyl)cyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4,4-dicarbomethoxycyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
20 5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine;
5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine;
25 5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4,4-diethylcyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4,4-di(hydroxymethyl)cyclopenten-1-yl]pyridine;
30 5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4,4-dicarbomethoxycyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
35 5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine;

- 4-[2-(5-fluoropyridin-2-yl)-4,4-dimethylcyclopenten-1-yl]
benzenesulfonamide;
- 4-[2-(5-fluoropyridin-2-yl)-4,4-diethylcyclopenten-1-yl]
benzenesulfonamide;
- 5 4-[2-(5-fluoropyridin-2-yl)-4,4-di(hydroxymethyl)
cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(5-fluoropyridin-2-yl)-4,4-dicarbomethoxycyclopenten-
1-yl]benzenesulfonamide;
- 4-[2-(5-fluoropyridin-2-yl)-4,4-difluorocyclopenten-1-yl]
10 benzenesulfonamide;
- 4-[2-(5-fluoropyridin-2-yl)-4,4-di(trifluoromethyl)
cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(5-fluoropyridin-2-yl)-4,4-di(fluoromethyl)
cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-
dimethylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-
diethylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-
20 di(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-
dicarbomethoxycyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-
difluorocyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-
di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-
di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 2-[2-phenylcyclopenten-1-yl]-5(methylsulfonyl)pyridine;
- 30 2-[2-(4-fluorophenyl)cyclopenten-1-yl]-
5(methylsulfonyl)pyridine;
- 2-[2-(4-chlorophenyl)cyclopenten-1-yl]-5-
(methylsulfonyl)pyridine;
- 2-[2-(4-methylphenyl)cyclopenten-1-yl]-5-
35 (methylsulfonyl)pyridine;
- 2-[2-(4-cyanophenyl)cyclopenten-1-yl]-
5(methylsulfonyl)pyridine;
- 2-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-5

- (methylsulfonyl)pyridine;
2-[2-phenylcyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
5 2-[2-(4-chlorophenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-methylphenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-cyanophenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
10 2-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)-4-methylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
15 2-[2-(4-fluorophenyl)-4-ethylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4-(hydroxymethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4-fluorocyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
20 2-[2-(4-fluorophenyl)-4-(trifluoromethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
25 2-[2-(4-fluorophenyl)-4-(fluoromethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4-methylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4-ethylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
30 2-[2-(4-trifluoromethylphenyl)-4-(hydroxymethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4-carbomethoxycyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
35 2-[2-(4-trifluoromethylphenyl)-4-fluorocyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-5-(methylsulfonyl)pyridine;


- 2-[2-(4-trifluoromethylphenyl)-4-(trifluoromethyl)
cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4-methylcyclopenten-1-yl]pyridine-5-
sulfonamide;
5 2-[2-(4-fluorophenyl)-4-ethylcyclopenten-1-yl]
pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)-4-(hydroxymethyl)cyclopenten-1-
yl]pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)-4-carbomethoxycyclopenten-1-yl]
10 pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)-4-fluorocyclopenten-1-yl]pyridine-5-
sulfonamide;
2-[2-(4-fluorophenyl)-4-(trifluoromethyl)cyclopenten-1-yl]
pyridine-5-sulfonamide;
15 2-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]pyridine-5-
sulfonamide;
2-[2-(4-fluorophenyl)-4-(fluoromethyl)cyclopenten-1-yl]
pyridine-5-sulfonamide;
2-[2-(4-trifluoromethylphenyl)-4-methylcyclopenten-1-
20 yl]pyridine-5-sulfonamide;
2-[2-(4-trifluoromethylphenyl)-4-ethylcyclopenten-1-yl]
pyridine-5-sulfonamide;
2-[2-(4-trifluoromethylphenyl)-4-(hydroxymethyl)
cyclopenten-1-yl]pyridine-5-sulfonamide;
25 2-[2-(4-trifluoromethylphenyl)-4-carbomethoxycyclopenten-1-
yl]pyridine-5-sulfonamide;
2-[2-(4-trifluoromethylphenyl)-4-fluorocyclopenten-1-yl]
pyridine-5-sulfonamide;
2-[2-(4-trifluoromethylphenyl)-4-(trifluoromethyl)
30 cyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]
pyridine-5-sulfonamide;
2-[2-(4-trifluoromethylphenyl)-4-(fluoromethyl)cyclopenten-
1-yl]pyridine-5-sulfonamide;
35 2-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-5-
(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-5-
(methylsulfonyl)pyridine;

- 2-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
5 2-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
10 2-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4,4-diethylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
15 2-[2-(4-trifluoromethylphenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
20 2-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-trifluoromethylphenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
25 2-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]pyridine-5-sulfonamide;
30 2-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]pyridine-5-sulfonamide;
2-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]pyridine-5-sulfonamide;
35 2-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine-5-sulfonamide;

- 2-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
 2-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]pyridine-5-sulfonamide;
 5 2-[2-(4-trifluoromethylphenyl)-4,4-diethylcyclopenten-1-yl]pyridine-5-sulfonamide;
 2-[2-(4-trifluoromethylphenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
 2-[2-(4-trifluoromethylphenyl)-4,4-
 10 dicarbomethoxycyclopenten-1-yl]pyridine-5-sulfonamide;
 2-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]pyridine-5-sulfonamide;
 2-[2-(4-trifluoromethylphenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine-5-sulfonamide; and
 15 2-[2-(4-trifluoromethylphenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine-5-sulfonamide.

Within Formula I there is a fourth subclass of compounds of high interest represented by Formula V:

20



V

- wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and
 25 wherein each of R³ through R⁷, R⁸, R¹⁰, R¹¹ and R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl
 30 and sulfamyl;
 or a pharmaceutically-acceptable salt thereof.

A preferred class of compounds consists of those compounds of Formula V wherein each of R^1 and R^2 is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxy carbonyl and carboxyl; and wherein each of R^3 through R^7 , R^8 , R^{10} , R^{11} and R^{12} is independently selected from hydrido, halo, lower alkyl, lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl.

A class of compounds of particular interest consists of those compounds of Formula V wherein each of R^1 and R^2 is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R^3 through R^8 and R^{10} through R^{12} is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl,

methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl.

5 A family of specific compounds of particular interest within Formula V consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 1-[2-(2,3-dimethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(3-fluoro-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(3-methoxy-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2,3-difluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(3-methyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(3-methoxy-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2,3-dichloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(3-methyl-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-methoxy-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(2,3-dimethoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(3-fluoro-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-[2,3-di(trifluoromethyl)pyridin-5-yl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(3-fluoro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(3-difluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(3-difluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-[2,3-di(difluoromethyl)pyridin-5-yl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(3-methyl-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-methoxy-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 4-[2-(2,3-dimethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-fluoro-2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 5 4-[2-(3-chloro-2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(2,3-difluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(3-chloro-2-fluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-fluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(2,3-dichloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(3-fluoro-2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 30 4-[2-(2,3-dimethoxypyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-(3-fluoro-2-methoxypyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(3-chloro-2-methoxypyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-[2,3-di(trifluoromethyl)pyridin-5-yl]cyclopenten-1-yl]benzenesulfonamide;
- 5 4-[2-(3-methyl-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-fluoro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(3-chloro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(3-difluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-[2,3-di(difluoromethyl)pyridin-5-yl]cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(3-fluoro-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-chloro-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 30 5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
- 2-chloro-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
- 35 2-methyl-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;

- 2-cyano-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
- 5 4-[2-(pyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-fluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-cyanopyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 15 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-methylcyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-ethylcyclopenten-1-yl]pyridine;
- 20 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-(hydroxymethyl)cyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-carbomethoxycyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-fluorocyclopenten-1-yl]pyridine;
- 25 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)cyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-carboxycyclopenten-1-yl]pyridine;
- 30 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-(fluoromethyl)cyclopenten-1-yl]pyridine;
- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-methylcyclopenten-1-yl]pyridine;
- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-ethylcyclopenten-1-yl]pyridine;
- 35 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-(hydroxymethyl)cyclopenten-1-yl]pyridine;

- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-carbomethoxycyclopenten-1-yl]pyridine;
2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-fluorocyclopenten-1-yl]pyridine;
5 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)cyclopenten-1-yl]pyridine;
2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-carboxycyclopenten-1-yl]pyridine;
2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-(fluoromethyl)cyclopenten-1-yl]pyridine;
10 4-[2-(2-fluoropyridin-5-yl)-4-methylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-fluoropyridin-5-yl)-4-ethylcyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(2-fluoropyridin-5-yl)-4-(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-fluoropyridin-5-yl)-4-carbomethoxycyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-fluoropyridin-5-yl)-4-fluorocyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(2-fluoropyridin-5-yl)-4-(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-fluoropyridin-5-yl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(2-fluoropyridin-5-yl)-4-(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine);
4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-diethylcyclopenten-1-yl]pyridine);
30 4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(hydroxymethyl)cyclopenten-1-yl]pyridine);
4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dicarbomethoxycyclopenten-1-yl]pyridine);
35 4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine);
4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine);

- 4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dicarboxycyclopenten-1-yl]pyridine;
- 4-[2-(2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-difluoromethylcyclopenten-1-yl]pyridine;
- 5 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-diethylcyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(hydroxymethyl)cyclopenten-1-yl]pyridine;
- 10 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dicarbomethoxycyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
- 15 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine;
- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine;
- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
- 20 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-diethylcyclopenten-1-yl]pyridine;
- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(hydroxymethyl)cyclopenten-1-yl]pyridine;
- 25 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dicarbomethoxycyclopenten-1-yl]pyridine;
- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
- 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine;
- 30 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine;
- 4-[2-(2-fluoropyridin-5-yl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-(2-fluoropyridin-5-yl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-fluoropyridin-5-yl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(2-fluoropyridin-5-yl)-4,4-dicarbomethoxycyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-fluoropyridin-5-yl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
- 5 4-[2-(2-fluoropyridin-5-yl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-fluoropyridin-5-yl)-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-dicarbomethoxycyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 5-(2-phenylcyclopenten-1-yl)-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 25 5-[2-(4-chlorophenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-methylphenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 30 5-[2-(4-cyanophenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-(2-phenylcyclopenten-1-yl)pyridine-2-sulfonamide;
- 35 5-[2-(4-fluorophenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-chlorophenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;

- 5-[2-(4-methylphenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-cyanophenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5 5-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4-methylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4-ethylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 10 5-[2-(4-fluorophenyl)-4-(hydroxymethyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4-carbomethoxycyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 15 5-[2-(4-fluorophenyl)-4-fluorocyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4-(trifluoromethyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 20 5-[2-(4-fluorophenyl)-4-(fluoromethyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4-methylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 25 5-[2-(4-trifluoromethylphenyl)-4-ethylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4-(hydroxymethyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4-carbomethoxycyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 30 5-[2-(4-trifluoromethylphenyl)-4-fluorocyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4-(trifluoromethyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 35 5-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4-(fluoromethyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;


- 5-[2-(4-fluorophenyl)-4-methylcyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4-ethylcyclopenten-1-yl]pyridine-2-sulfonamide;
- 5 5-[2-(4-fluorophenyl)-4-(hydroxymethyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4-carbomethoxycyclopenten-1-yl]pyridine-2-sulfonamide;
- 10 5-[2-(4-fluorophenyl)-4-fluorocyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4-(trifluoromethyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]pyridine-2-sulfonamide;
- 15 5-[2-(4-fluorophenyl)-4-(fluoromethyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-trifluoromethylphenyl)-4-methylcyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-trifluoromethylphenyl)-4-ethylcyclopenten-1-yl]pyridine-2-sulfonamide;
- 20 5-[2-(4-trifluoromethylphenyl)-4-(hydroxymethyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-trifluoromethylphenyl)-4-carbomethoxycyclopenten-1-yl]pyridine-2-sulfonamide;
- 25 5-[2-(4-trifluoromethylphenyl)-4-fluorocyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-trifluoromethylphenyl)-4-(trifluoromethyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]pyridine-2-sulfonamide;
- 30 5-[2-(4-trifluoromethylphenyl)-4-(fluoromethyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 35 5-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;

- 5-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-
2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-2-
(methylsulfonyl)pyridine;
- 5 5-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-
yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-
yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-
10 yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4,4-diethylcyclopenten-1-
yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4,4-di(hydroxymethyl)
cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 15 5-[2-(4-trifluoromethylphenyl)-4,4-
dicarbomethoxycyclopenten-1-yl]-2-(methylsulfonyl)
pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-
yl]-2-(methylsulfonyl)pyridine;
- 20 5-[2-(4-trifluoromethylphenyl)-4,4-di(trifluoromethyl)
cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4,4-di(fluoromethyl)
cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-
25 yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]pyridine-
2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-
yl]pyridine-2-sulfonamide;
- 30 5-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-
yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-
yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-
35 yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-
yl]pyridine-2-sulfonamide;

5-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]pyridine-2-sulfonamide;
 5-[2-(4-trifluoromethylphenyl)-4,4-diethylcyclopenten-1-yl]pyridine-2-sulfonamide;
 5 5-[2-(4-trifluoromethylphenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
 5-[2-(4-trifluoromethylphenyl)-4,4-dicarbomethoxycyclopenten-1-yl]pyridine-2-sulfonamide;
 5-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]pyridine-2-sulfonamide;
 10 5-[2-(4-trifluoromethylphenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]pyridine-2-sulfonamide; and
 5-[2-(4-trifluoromethylphenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine-2-sulfonamide.

15

Within Formula I there is a fifth subclass of compounds of high interest represented by Formula VI:



VI

20

wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl;

wherein R⁵ is selected from alkylsulfonyl, haloalkylsulfonyl and sulfamyl; and

25 wherein each of R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹¹ and R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl and alkoxyalkyl; or a pharmaceutically-
 30 acceptable salt thereof.

A preferred class of compounds consists of those compounds of Formula VI wherein each of R^1 and R^2 is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxy carbonyl and carboxyl; wherein R^5 is selected from lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl; and wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^9 , R^{11} and R^{12} is independently selected from hydrido, halo, lower alkyl, lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower haloalkoxy, lower hydroxyalkyl and lower alkoxyalkyl.

A class of compounds of particular interest consists of those compounds of Formula VI wherein each of R^1 and R^2 is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; wherein R^5 is selected from methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl; and wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^9 , R^{11} and R^{12} is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl,

methanesulfonyl, ethanesulfonyl, trifluoromethanesulfonyl and sulfamyl.

A family of specific compounds of particular
5 interest within Formula VI consists of compounds and
pharmaceutically-acceptable salts thereof as follows:

- 4-[2-[4-(methanesulfonyl)phenyl]cyclopenten-1-yl]pyridine;
4-[2-(4-pyridinyl)cyclopenten-1-yl]benzenesulfonamide;
10 4-[2-[4-(methanesulfonyl)phenyl]-4,4-dimethylcyclopenten-1-
yl]pyridine;
4-[2-[4-(methanesulfonyl)phenyl]-4,4-diethylcyclopenten-1-
yl]pyridine;
4-[2-[4-(methanesulfonyl)phenyl]-4,4-
15 di(hydroxymethyl)cyclopenten-1-yl]pyridine;
4-[2-[4-(methanesulfonyl)phenyl]-4,4-
dicarbomethoxycyclopenten-1-yl]pyridine;
4-[2-[4-(methanesulfonyl)phenyl]-4,4-difluorocyclopenten-1-
yl]pyridine;
20 4-[2-[4-(methanesulfonyl)phenyl]-4,4-
di(trifluoromethyl)cyclopenten-1-yl]pyridine;
4-[2-[4-(methanesulfonyl)phenyl]-4,4-
di(fluoromethyl)cyclopenten-1-yl]pyridine;
4-[2-[4-(pyridinyl)phenyl]-4,4-dimethylcyclopenten-1-
25 yl]benzenesulfonamide;
4-[2-[4-(pyridinyl)phenyl]-4,4-diethylcyclopenten-1-
yl]benzenesulfonamide;
4-[2-[4-(pyridinyl)phenyl]-4,4-di(hydroxymethyl)
cyclopenten-1-yl]benzenesulfonamide;
30 4-[2-[4-(pyridinyl)phenyl]-4,4-dicarbomethoxycyclopenten-1-
yl]benzenesulfonamide;
4-[2-[4-(pyridinyl)phenyl]-4,4-difluorocyclopenten-1-
yl]benzenesulfonamide;
4-[2-[4-(pyridinyl)phenyl]-4,4-di(trifluoromethyl)
35 cyclopenten-1-yl]benzenesulfonamide; and
4-[2-[4-(pyridinyl)phenyl]-4,4-di(fluoromethyl)cyclopenten-
1-yl]benzenesulfonamide.

Where used, either alone or within other terms such as "haloalkyl", "alkoxyalkyl" and "hydroxyalkyl", the term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about five carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl and the like. The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH₂-) radical. The term "halo" means halogens such as fluorine, chlorine, bromine or iodine. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have either a bromo, chloro or a fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. More preferred haloalkyl radicals are "lower haloalkyl" radicals having one to six carbon atoms. Examples of such lower haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms. Examples of such lower hydroxyalkyl radicals include hydroxymethyl, 2-

hydroxyethyl, 4-hydroxybutyl and 6-hydroxyhexyl. The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals are

5 "lower alkoxy" radicals having one to six carbon atoms. Examples of such lower alkoxy radicals include methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy and methylenedioxy. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy radicals attached to the

10 alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals. More preferred alkoxyalkyl radicals are "lower alkoxyalkyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkoxyalkyl radicals include methoxymethyl, methoxyethyl and the like.

15 The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy or haloalkoxyalkyl radicals. The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to about ten

20 carbon atoms attached to a divalent sulfur atom, such as a methylthio radical. More preferred alkylthio radicals are "lower alkylthio" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthio radicals include methylthio, ethylthio, propylthio, butylthio and

25 hexylthio. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl and biphenyl. Preferred aryl radicals are those consisting of one, two, or three benzene rings. The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent

30 radicals $-SO_2-$. "Alkylsulfonyl" embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylsulfonyl radicals

35 include methylsulfonyl and ethylsulfonyl. The term "haloalkylsulfonyl" embraces haloalkyl radicals attached to a sulfonyl radical, where haloalkyl and alkyl are defined as above. More preferred haloalkylsulfonyl radicals are

"lower haloalkylsulfonyl" radicals having lower haloalkyl radicals as described above. Examples of such lower haloalkylsulfonyl radicals include fluoromethylsulfonyl and trifluoromethylsulfonyl. The terms "sulfamyl" or

5 "sulfonamidyl" denotes a sulfonyl radical substituted with an amine radical, forming a sulfonamide ($-\text{SO}_2\text{NH}_2$). The term "carboxyl", whether used alone or with other terms, denotes $-\text{CO}_2\text{H}$. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached

10 via a carbon atom to a "carbonyl" radical ($\text{C}=\text{O}$). More preferred alkoxycarbonyl radicals are "lower alkoxycarbonyl" radicals having lower alkoxy radicals as described above. Examples of such "lower alkoxycarbonyl" radicals include methoxycarbonyl, ethoxycarbonyl,

15 isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and tert-butoxycarbonyl ($(\text{CH}_3)_3\text{CO}_2\text{C}-$).

The present invention comprises a pharmaceutical

20 composition for the treatment of inflammation and inflammation-associated disorders, such as arthritis, comprising a therapeutically-effective amount of a compound of Formula I in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent.

25

The present invention also comprises a therapeutic method of treating inflammation or inflammation-associated disorders in a subject, the method comprising administering to a subject having such

30 inflammation or disorder a therapeutically-effective amount of a compound of Formula I.

Also included in the family of compounds of Formula I are isomeric forms including diastereoisomers and


35 the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not

critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, 2-hydroxyethanesulfonic, toluenesulfonic, sulfanilic, cyclohexylaminosulfonic, stearic, algenic, β -hydroxybutyric, salicylic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I.

GENERAL SYNTHETIC PROCEDURES


The compounds of the invention can be synthesized according to the following procedures of Schemes I-XXII, wherein the R¹-R¹² substituents are as defined for Formula I, above, except where further noted.

Scheme I




Synthetic Scheme I shows the preparation of 1,2-dibromocyclopentene (3) in two steps from commercially available cyclopentanone (1) using a procedure similar to the one developed by Montgomery, et al., [*J.Am.Chem.Soc.*, 87, 1917 (1965)]. In step one, chlorination with phosphorus pentachloride gives 1-chlorocyclopentene (2). In step two, bromination of 2, followed by the elimination of hydrogen chloride on treatment with potassium tert-butoxide, provides 3.

Scheme II




Synthetic Scheme II shows the preparation of 1,2-dibromo-4,4-dicarbomethoxycyclopentene (8) in four steps from commercially available mucobromic acid (4) using a procedure similar to the one developed by Lerstrub, et al., [*Syn.Metals*, 19, 647 (1987)]. In step one, mucobromic acid is converted to its ethyl ester 5 on treatment with ethanol in toluene at reflux in the presence of p-toluenesulfonic acid (TsOH). In step two, reduction of 5 with diisobutylaluminum hydride (DIBAL) in methylene chloride gives the diol 6. In step three, the diol 6 is reacted with thionyl chloride in dimethylformamide (DMF) to give the corresponding dichloride 7. In step four, the dichloride 7 is dialkylated with the dianion of methyl malonate to give 8.

Scheme III




Synthetic Scheme III shows the preparation of 1,2-dibromo-4,4-dicarboxycyclopentene (9) and 1,2-bromo-4,4-di(hydroxymethyl)cyclopentene (10) from synthetic intermediate 8 (prepared in Synthetic Scheme II). Reaction of 8 with lithium hydroxide in tetrahydrofuran (THF) followed by careful acidification at 0°C gives the diacid 9; treatment with DIBAL gives the corresponding diol 10.

Scheme IV




Synthetic Scheme IV shows the preparation of 1,2-dibromo-4-carboxycyclopentene (11) and 1,2-dibromo-4,4-bis(trifluoromethyl)cyclopentene (12) from synthetic intermediate 9 (prepared in Synthetic Scheme III). On heating, the diacid 9 is converted to the monoacid 11; treatment with selenium tetrafluoride in 1,1,2-trichlorotrifluoroethane at reflux gives the bistrifluoromethyl analog 12.

Scheme V




Synthetic Scheme V shows the preparation of 1,2-dibromo-4,4-di(fluoromethyl)cyclopentene (13) and 1,2-dibromo-4,4-dimethylcyclopentene (16) from synthetic intermediate 10 (prepared in Synthetic Scheme III). Treatment of the diol 10 with diethylaminosulfurtrifluoride (DAST) in methylene chloride gives the corresponding fluoromethyl analog 13. Reaction of 10 with p-toluenesulfonyl chloride (TsCl) in the presence of pyridine gives the ditosylate 14. Reaction of 14 with sodium iodide in DMF/water (3:1) at 150°C gives the di(iodomethyl) analog 15 which is subsequently reduced with sodium cyanoborohydride to give the dimethyl analog 16.

Scheme VI




Synthetic Scheme VI shows the preparation of 1,2-dibromo-4,4-difluorocyclopentene (**19**) from synthetic intermediate **7** (prepared in Synthetic Scheme II). Using a procedure similar to the one developed by Ogura, et al., [*Tetrahedron Lett.*, **32**, 2767 (1975)], the dianion of methyl methylthiomethyl sulfoxide (generated by potassium hydride in THF) is reacted with **7** to give the dimethyl dithioacetal S-oxide **17**. Subsequent hydrolysis with 9 N sulfuric acid gives the corresponding ketone **18**. Reaction of **18** with selenium tetrafluoride in 1,1,2-trichlorotrifluoroethane at reflux gives the 4,4-difluoro analog **19**.

Scheme VII



5 Synthetic Scheme VII shows the preparation
of the prerequisite substituted phenylboronic acids 21
and substituted pyridinylboronic acids 23, 25, and 27
from the available bromides 20, 22, 24, and 26,
respectively. Halogen-metal interchange in THF at
10 -78°C generates the corresponding organolithium
reagents which are reacted with trimethyl borate.
Hydrolysis with 3N hydrochloric acid provides the
substituted phenylboronic acids 21 and the substituted
pyridinylboronic acids 23, 25, and 27, respectively.

Scheme VIII




5

Synthetic Scheme VIII shows the preparation of the prerequisite substitutedphenyltrimethyltin analogs **29** and substitutedpyridinyltrimethyltin analogs **31**, **33**, and **35** from the available bromides **28**, **30**, **32**, and **34**, respectively. Halogen-metal interchange in THF at -78°C generates the organolithium reagents which are reacted with trimethyltin chloride. Purification provides the substitutedphenyltrimethyltin analogs **29** and the substitutedpyridinyltrimethyltin analogs **31**, **33**, and

15

35, respectively.

Scheme IX




5

Synthetic Scheme IX shows the four step procedure for the preparation of 1,2-diaryl-4,4-disubstitutedcyclopentene anti-inflammatory agents **41** from the available bromides **20** using a sequential coupling procedure which is similar to the coupling procedure developed by Negishi, et al., [*J.Org.Chem.*, **42**, 1821 (1977)]. In step one, halogen-metal

interchange of 20 with n-butyllithium in THF at -78°C gives the corresponding organolithium reagents which subsequently react with zinc chloride to give the organozinc reagents 36. In step two, the organozinc reagents 36 are coupled with the 1,2-dibromo-4,4-disubstitutedcyclopentenenes 37 (prepared in Synthetic Schemes I-IV) in the presence of a Pd⁰ catalyst, e.g., tetrakis(triphenylphosphine)palladium (0), to give the monocoupled bromides 38 (after separation from the biscoupled by-product). In step three, the bromides 38 are treated as above to give the organozinc reagents 39. In step four, the monocoupled organozinc reagents 39 are coupled with the arylbromides 40 (which can be identical with 20 when R³=R⁸, R⁴=R⁹, R⁵=R¹⁰, R⁶=R¹¹, and R⁷=R¹²) to give the 1,2-diaryl-4,4-disubstitutedcyclopentene anti-inflammatory agents 41 of this invention.

Scheme X




- 5 Synthetic Scheme X shows the two step
 procedure for the preparation of 1,2-diaryl-4,4-
 disubstitutedcyclopentene anti-inflammatory agents **41**
 from 1,2-dibromo-4,4-disubstitutedcyclopentenenes **37**
 (prepared in Synthetic Scheme I-VI) and substituted
 10 phenylboronic acids **21** and **42** (prepared in Synthetic
 Scheme VII) using a sequential coupling procedure
 which is similar to the coupling procedure developed
 by Suzuki, et al., [*Syn. Commun.*, **11**, 513 (1981)]. In
 step one, the dibromides **37** are treated with the
 15 boronic acids **21** in toluene at reflux in the presence
 of Pd⁰ catalyst, e.g.,
 tetrakis(triphenylphosphine)palladium (0), and 2M

sodium carbonate to give the monocoupled bromides **38** (after separation from the biscoupled by-product). In step two, the monocoupled bromides **38** are coupled as above with the boronic acids **42** (which can be identical with **21** when $R^3=R^8$, $R^4=R^9$, $R^5=R^{10}$, $R^6=R^{11}$, and $R^7=R^{12}$) give the 1,2-diaryl-4,4-disubstitutedcyclopentene anti-inflammatory agents **41** of this invention.

10

Scheme XI




Synthetic Scheme XI shows the two step procedure for the preparation of 1,2-diaryl-4,4-disubstitutedcyclopentene anti-inflammatory agents **41** from 1,2-dibromo-4,4-disubstitutedcyclopentenenes **37** (prepared in Synthetic Scheme I-IV) and substitutedphenyltrimethyltin analogs **29** and **43** (prepared in Synthetic Scheme VII) using a sequential

20

coupling procedure which is similar to the coupling procedure developed by Stille, et al., [J.Am.Chem.Soc., 101, 4992 (1979)]. In step one, the dibromides **37** are treated with the trimethyltin analogs **29** in toluene at reflux in the presence of a Pd⁰ catalyst, e.g., tetrakis(triphenylphosphine)palladium (0), to give the monocoupled bromides **38** (after separation from the biscoupled by-product). In step two, the monocoupled bromides **38** are coupled as above with the trimethyltin analogs **43** (which can be identical with **29** when R³=R⁸, R⁴=R⁹, R⁵=R¹⁰, R⁶=R¹¹, and R⁷=R¹²) to give the 1,2-diaryl-4,4-disubstitutedcyclopentene anti-inflammatory agents **41** of this invention.


Scheme XII



Synthetic Scheme XII shows the three step preparation of 1-aryl-2-(2-pyridinyl)-4,4-disubstitutedcyclopentene anti-inflammatory agents **46** from 1,2-dibromo-4,4-disubstitutedcyclopentenenes **37** (prepared in Synthetic Scheme I-IV) and the available

2-bromopyridines **22**. In step one, halogen-metal interchange of **22** with n-butyllithium in THF at -78°C gives the 2-lithiopyridines which subsequently react with zinc chloride to give the corresponding 2-pyridinylzinc reagents **44**. In step two, a Negishi coupling (see Synthetic Scheme IX) of the 2-pyridinylzinc reagents **44** with **37** gives the monocoupled 2-pyridinyl bromides **45** (after separation from the biscoupled by-product). In step three, a Suzuki coupling (see Synthetic Scheme X) of the monocoupled 2-pyridinyl bromides **45** with substituted phenylboronic acids **21** (prepared in Synthetic Scheme VII) gives the 1-aryl-2-(2-pyridinyl)-4,4-disubstitutedcyclopentene anti-inflammatory agents **46** of this invention.


Scheme XIII



Synthetic Scheme XIII shows the three step preparation of 1-aryl-2-(3-pyridinyl)-4,4-disubstitutedcyclopentene anti-inflammatory agents **49** from 1,2-dibromo-4,4-disubstitutedcyclopentenenes **37**

(prepared in Synthetic Scheme I-IV) and the available 3-bromopyridines **24**. In step one, halogen-metal interchange of **24** with n-butyllithium in THF at -78°C gives the 3-lithiopyridines which subsequently react with zinc chloride to give the corresponding 3-pyridinylzinc reagents **47**. In step two, a Negishi coupling (see Synthetic Scheme IX) of the 3-pyridinylzinc reagents **47** with **37** gives the monocoupled 3-pyridinyl bromides **48** (after separation from the biscoupled by-product). In step three, a Suzuki coupling (see Synthetic Scheme X) of the monocoupled 3-pyridinyl bromides **48** with substituted phenylboronic acid **21** (prepared in Synthetic Scheme VII) gives the 1-aryl-2-(3-pyridinyl)-4,4-disubstitutedcyclopentene anti-inflammatory agents **49** of this invention.


Scheme XIV




Synthetic Scheme XIV shows the three step preparation of 1-aryl-2-(4-pyridinyl)-4,4-disubstitutedcyclopentene anti-inflammatory agents **52**

from 1,2-dibromo-4,4-disubstitutedcyclopentenenes **37**
(prepared in Synthetic Scheme I-IV) and the available
4-bromopyridines **26**. In step one, halogen-metal
interchange of **26** with n-butyllithium in THF at -78°C
5 gives the 4-lithiopyridines which subsequently react
with zinc chloride to give the corresponding 4-
pyridinylzinc reagents **50**. In step two, a Negishi
coupling (see Synthetic Scheme IX) of the 4-
pyridinylzinc reagents **50** with **37** gives the
10 monocoupled 4-pyridinyl bromides **51** (after separation
from the biscoupled by-product). In step three, a
Suzuki coupling (see Synthetic X) of the monocoupled
4-pyridinyl bromides **51** with substituted phenylboronic
acids **21** (prepared in Synthetic Scheme VII) gives the
15 1-aryl-2-(4-pyridinyl)-4,4-disubstitutedcyclopentene
anti-inflammatory agents **52** of this invention.


Scheme XV

**53**


1. $\text{Cl-CO-OCH}_2\text{CH}(\text{CH}_3)_2$, THF,
0 °C, $\text{N}(\text{C}_2\text{H}_5)_3$
2. $\text{HN}(\text{CH}_3)\text{OCH}_3$, HCl, $\text{N}(\text{C}_2\text{H}_5)_3$

**54**

CH_3MgBr ,
THF, 0 °C


**55**

$(\text{CH}_3)_3\text{Si-Cl}$, NaI,
 $\text{N}(\text{C}_2\text{H}_5)_3$, CH_3CN

**56**

Synthetic Scheme XV shows the three step procedure used to prepare the phenyl silyl enol ethers 56 from commercially available benzoic acids 53. In
5 step one, a THF solution at 0°C of the benzoic acids 53 and two equivalents of triethylamine are sequentially treated with isobutyl chloroformate and N-hydroxymethyl-N-methylamine hydrochloride to give the Weinreb amides 54 [see: S. Nahm and S.M. Weinreb,
10 *Tetrahedron Lett.*, 21, 3815 (1981)]. In step two, the amides 54 are reacted with methylmagnesium bromide to give the corresponding acetophenones 55. In step three, the acetophenones 55 are treated with chlorotrimethylsilane in acetonitrile in the presence
15 of triethylamine and sodium iodide to give the corresponding phenyl silyl enol ethers. 56.


Scheme XVI



57




1. $Cl-CO-OCH_2CH(CH_3)_2$, THF, $0^\circ C$, $N(C_2H_5)_3$
2. $HN(CH_3)OCH_3$ HCl, $N(C_2H_5)_3$



58




CH_3MgBr ,
THF, $0^\circ C$



59




$(CH_3)_3Si-Cl$, NaI,
 $N(C_2H_5)_3$, CH_3CN



60


Synthetic Scheme XVI shows the three step procedure used to prepare the phenyl silyl enol ethers 60 from commercially available benzoic acids 57. In
5 step one, a THF solution at 0°C of the benzoic acids 57 and two equivalents of triethylamine are sequentially treated with isobutyl chloroformate and N-hydroxymethyl-N-methylamine hydrochloride to give the Weinreb amides 58. In step two, the amides 58 are
10 reacted with methylmagnesium bromide to give the corresponding acetophenones 59. In step three, the acetophenones 59 are treated with chlorotrimethylsilane in acetonitrile in the presence of triethylamine and sodium iodide to give the
15 corresponding phenyl silyl enol ethers 60.

Scheme XVII




61

1. $Cl-CO-OCH_2CH(CH_3)_2$, THF,
0 °C, $N(C_2H_5)_3$
2. $HN(CH_3)OCH_3$, HCl, $N(C_2H_5)_3$




62

- CH_3MgBr ,
THF, 0 °C



63


- $(CH_3)_3Si-Cl$, NaI,
 $N(C_2H_5)_3$, CH_3CN




64

Synthetic Scheme XVII shows the three step procedure used to prepare the 2-pyridinyl silyl enol ethers **64** from commercially available picolinic acids **61**. In step one, a THF solution at 0°C of the picolinic acids **61** and two equivalents of triethylamine are sequentially treated with isobutyl chloroformate and N-hydroxymethyl-N-methylamine hydrochloride to give the Weinreb amides **62**. In step two, the amides **62** are reacted with methylmagnesium bromide to give the corresponding 2-acetylpyridines **63**. In step three, the 2-acetylpyridines **63** are treated with chlorotrimethylsilane in acetonitrile in the presence of triethylamine and sodium iodide to give the corresponding 2-pyridinyl silyl enol ethers **64**.


Scheme XVIII

**65**


1. $\text{Cl}-\text{CO}-\text{OCH}_2\text{CH}(\text{CH}_3)_2$, THF, 0°C , $\text{N}(\text{C}_2\text{H}_5)_3$
2. $\text{HN}(\text{CH}_3)\text{OCH}_3 \text{ HCl}$, $\text{N}(\text{C}_2\text{H}_5)_3$

**66**

CH_3MgBr ,
THF, 0°C


**67**

$(\text{CH}_3)_3\text{Si}-\text{Cl}$, NaI ,
 $\text{N}(\text{C}_2\text{H}_5)_3$, CH_3CN

**68**


Synthetic Scheme XVIII shows the three step procedure used to prepare the 3-pyridinyl silyl enol ethers 68 from commercially available nicotinic acids 65. In step one, a THF solution at 0°C of the nicotinic acids 65 and two equivalents of triethylamine are sequentially treated with isobutyl chloroformate and N-hydroxymethyl-N-methylamine hydrochloride to give the Weinreb amides 66. In step two, the amides 66 are reacted with methylmagnesium bromide to give the corresponding 3-acetylpyridines 67. In step three, the 3-acetylpyridines 67 are treated with chlorotrimethylsilane in acetonitrile in the presence of triethylamine and sodium iodide to give the corresponding 3-pyridinyl silyl enol ethers 68

Scheme XIX




69

1. Cl-CO-OCH₂CH(CH₃)₂, THF,
0 °C, N(C₂H₅)₃
2. HN(CH₃)OCH₃ HCl, N(C₂H₅)₃




70

CH₃MgBr,
THF, 0 °C



71


(CH₃)₃Si-Cl, NaI,
N(C₂H₅)₃, CH₃CN



72

Synthetic Scheme XIX shows the three step procedure used to prepare the 4-pyridinyl silyl enol ethers **72** from commercially available isonicotinic acids **69**. In step one, a THF solution at 0°C of the isonicotinic acids **69** and two equivalents of triethylamine are sequentially treated with isobutyl chloroformate and N-hydroxymethyl-N-methylamine hydrochloride to give the Weinreb amides **70**. In step two, the amides **70** are reacted with methylmagnesium bromide to give the corresponding 4-acetylpyridines **71**. In step three, the 4-acetylpyridines **71** are treated with chlorotrimethylsilane in acetonitrile in the presence of triethylamine and sodium iodide to give the corresponding 4-pyridinyl silyl enol ethers **72**.

Scheme XX




Synthetic Scheme XX shows the two step procedures used to prepare the enones **74**, **76**, **78**, and **80** from the phenyl silyl enol ethers **56**, **60**, **68**, and **72**, respectively (prepared in Synthetic Schemes XVI-XIX). In step one, the silyl enol ethers **56**, **60**, **68**, and **72** are reacted with the appropriate ketones in methylene chloride in the presence of titanium (IV) chloride to give the corresponding 3-hydroxyketones **73**, **75**, **77**, and **79**, respectively. In step two, the 3-hydroxyketones **73**, **75**, **77**, and **79** are dehydrated with trifluoroacetic anhydride and triethylamine in methylene chloride at $0^\circ C$ to give the corresponding

78

enones **74**, **76**, **78**, and **80**, respectively.


Scheme XXI



5 Synthetic Scheme XXI shows the procedures
 used to prepare the 1,5-diketones **81**, **82**, **83**, and **84**
 from the enones **74**, **76**, **78**, and **80**, respectively
 (prepared in Synthetic Scheme XX). Methylene chloride
 solutions of the enones **74**, **76**, **78**, and **80** are reacted
 10 with the silyl enol ethers **56** (prepared in Synthetic
 Scheme XV) in the presence of titanium (IV) chloride

to give the corresponding 1,5-diketones **81**, **82**, **83**, and **84**, respectively.

Scheme XXII




5

Synthetic Scheme XXII shows alternative procedures which may be used to prepare 1,2-diaryl-4,4-disubstitutedcyclopentenones **41**, 1-aryl-2-(2-

pyridinyl)-4,4-disubstitutedcyclopentenes **46**, 1-aryl-2-(3-pyridinyl)-4,4-disubstitutedcyclopentenes **49**, and 1-aryl-2-(4-pyridinyl)-4,4-disubstitutedcyclopentenes **52** from the 1,5-diketones **81**, **82**, **83** and **84**,
 5 respectively (prepared in Synthetic Scheme XXI).
 Tetrahydrofuran solutions of 1,5-diketones **81**, **82**, **83** and **84** are treated with metallic zinc and titanium IV) chloride to give the reduced cyclized 1,2-diaryl-4,4-disubstitutedcyclopentene antiinflammatory agents **41**,
 10 1-aryl-2-(2-pyridinyl)-4,4-disubstitutedcyclopentene antiinflammatory agents **46**, 1-aryl-2-(3-pyridinyl)-4,4-disubstituted cyclopentene antiinflammatory agents **49**, and 1-aryl-2-(4-pyridinyl)-4,4-disubstitutedcyclopentene antiinflammatory agents **52**,
 15 respectively, of this invention.

Scheme XXIII




Synthetic Scheme XXIII shows the three step
 20 procedure used to prepare sulfonamide antiinflammatory

agents 86 and the two step procedure used to prepare fluoromethyl sulfone antiinflammatory agents 87 from their corresponding methyl sulfones 85. In step one, THF solutions of the methyl sulfones 85 at -78°C are
5 treated with an alkylolithium reagent, e.g., methyllithium, n-butyllithium, etc. In step two, the anions generated in step one are treated with an organoborane, e.g., triethylborane, tributylborane, etc., at -78°C then allowed to warm to ambient
10 temperature prior at stirring at reflux. In step three, an aqueous solution of sodium acetate and hydroxyamine-O-sulfonic acid is added to provide the corresponding sulfonamide antiinflammatory agents 86 of this invention. Alternatively, the anion solutions
15 generated in step one may be warmed to 0°C and treated with N-fluorodibenzenesulfonamide to provide the corresponding fluoromethyl sulfone antiinflammatory agents 87 of this invention.

The following examples contain detailed
20 descriptions of the methods of preparation of compounds of Formula I-VI. These detailed descriptions fall within the scope, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These detailed descriptions are presented for
25 illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated.

Exempl 1



1-[2-(4-Fluorophenyl)cyclopenten-1-yl]-
4-(methylsulfonyl)benzene

5

Step 1: Preparation of 4-methylthiophenylboronic acid.

Under nitrogen, a stirred solution of 30 g
10 (150 mmol) of 4-bromothiobenzene (Aldrich) in 1500 mL of
anhydrous THF at -78°C was treated with 180 mmol of *n*-
butyllithium in hexane. After 30 minutes, 51 mL (450
mmol) of trimethylborate was added neat and the reaction
was allowed to warm to ambient temperature overnight. A
15 solution of 300 mL of 10% NaOH was added and the mixture
stirred vigorously for 1 hour. The solvent was removed
in vacuo, the pH adjusted to 4-5, and the product
collected by filtration. Repeated washings with hexane
and water provided 21 g (83%) of 4-
20 methylthiophenylboronic acid (21 in Synthetic Scheme VII
when $\text{R}^5 = \text{SCH}_3$ and $\text{R}^3, \text{R}^4, \text{R}^6$, and $\text{R}^7 = \text{H}$) as a colorless
solid: NMR ($\text{DMSO}-d_6$) δ 2.47 (s, 3H), 7.20 (d, $J = 8$ Hz,
2H), 7.71 (d, $J = 8$ Hz, 2H), 7.96 (br s, 2H).

25 Step 2: Preparation of 1-(2-bromocyclopenten-1-yl)-4-
(methylthio)benzene.

Under nitrogen, 36.4 g (161 mmol) of 1,2-
dibromocyclopentene (Aldrich) was reacted with 18.0 g
30 (107 mmol) of 4-methylthiophenylboronic acid (Step 1) in

550 mL of toluene, 365 mL of ethanol, and 235 mL of 2M Na₂CO₃ in the presence of 6.0 g (5 mol %) of Pd(PPh₃)₄. The reaction was vigorously stirred at reflux overnight and concentrated in vacuo. The residue was dissolved in ethyl acetate and washed with water, dried (MgSO₄), and reconcentrated. Purification by silica gel chromatography (Waters Prep-500) with hexane gave 9.39 g (22%) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (38 in Synthetic Scheme X when R⁵ = SCH₃ and R¹, R², R³, R⁴, R⁶, and R⁷ = H) as a solid: mp 52-54°C; NMR (CDCl₃) δ 1.98-2.09 (m, 2H), 2.50 (s, 3H), 2.70-2.78 (m, 2H), 2.80-2.89 (m, 2H), 7.24 (d, J = 8 Hz, 2H), 7.55 (d, J = 8 Hz, 2H).

15 Step 3: Preparation of 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene.


Under nitrogen, 1.5 g (5.6 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Step 2) was reacted with 1.5 g (11 mmol) of 4-fluorophenylboronic acid (Lancaster) in 30 mL of toluene, 20 mL of ethanol, and 25 mL of 2M Na₂CO₃ in the presence of 250 mg of Pd(PPh₃)₄. The reaction was vigorously stirred at reflux overnight and concentrated in vacuo. The residue was dissolved in ethyl acetate and washed with water, dried (MgSO₄), and reconcentrated. Purification by silica gel chromatography with hexane gave 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene (41 in Synthetic Scheme X when R⁵ = SCH₃, R¹⁰ = F, and R¹, R², R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹¹, and R¹² = H) as a colorless solid: mp 46-47°C; NMR (CDCl₃) δ 2.04 (m, J = 7 Hz, 2H), 2.45 (s, 3H), 2.86 (t, J = 7 Hz, 4H), 6.86-6.94 (m, 2H), 7.08 (br s, 4H), 7.10-7.18 (m, 2H).

35

Step 4: Preparation of 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

A solution of 1.5 g (5 mmol) of 1-[2-(4-fluorophenyl) cyclopenten-1-yl]-4-(methylthio)benzene (Step 3) in 46 mL of methanol/THF (1:1) was slowly treated with 5.2 g (8.4 mmol) of Oxone® [2 KHSO₅.KHSO₄.K₂SO₄] in 23 mL of water. After stirring for 4 hours, the solvent was removed in vacuo. The residue was dissolved in ethyl acetate and washed with water and brine, dried (MgSO₄), and reconcentrated. Recrystallization from ethyl acetate/hexane provided 960 mg (54%) of 1-[2-(4-fluorophenyl) cyclopenten-1-yl]-4-(methylsulfonyl)benzene (**41** in Synthetic Scheme X when R⁵= SO₂CH₃, R¹⁰= F, and R¹, R², R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹¹, and R¹²= H) as a colorless solid: mp 138-139°C; NMR (CDCl₃) δ 2.09 (m, J= 7 Hz, 2H), 2.91 (t, J= 7 Hz, 4H), 3.04 (s, 3H), 6.88-6.96 (m, 2H), 7.06-7.14 (m, 2H), 7.32 (d, J= 8 Hz, 2H), 7.76 (d, J= 8 Hz, 2H). MS (EI) m/e (rel intensity) 316 (100), 237 (41), 161 (13); HRMS. Calc'd for C₁₈H₁₇FO₂S: 316.0933. Found: 316.0943. Anal. Calc'd for C₁₈H₁₇FO₂S: C, 68.33; H, 5.42; F, 6.00; S, 10.13. Found: C, 68.08; H, 5.45; F, 6.42; S, 9.98.

Example 2



1-[2-(4-Fluoro-2-methylphenyl)cyclopenten-1-yl]-
4-(methylsulfonyl)benzene

Step 1: Preparation of 4-fluoro-2-methylphenylboronic acid.

Following the synthetic procedure outlined in Step 1 of Example 1, 2-bromo-5-fluorotoluene (Aldrich) was converted to 4-fluoro-2-methylphenylboronic acid:

5 NMR (DMSO-d₆) δ 2.40 (s, 3H), 6.85-6.99 (m, 2H), 7.46 (d, J = 7 Hz, 1H).

Step 2: Preparation of 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene.

10

Following the synthetic procedure outlined in Step 3 of Example 1, 500 mg (1.9 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1, 15 Step 2) was reacted with 590 mg (3.8 mmol) of 4-fluoro-2-methylphenylboronic acid (Step 1). Purification by silica gel chromatography (MPLC) with hexane gave 500 mg (95%) of 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene as a colorless solid: mp 67-20 68°C; NMR (CDCl₃) δ 2.00-2.11 (m, 2H), 2.05 (s, 3H), 2.41 (s, 3H), 2.69-2.77 (m, 2H), 2.86-2.95 (m, 2H), 6.80-7.07 (m, 7H).


Step 3: Preparation of 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

25

Following the synthetic procedure outlined in Step 4 of Example 1, 470 mg (1.6 mmol) of 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene (Step 2) was oxidized. Purification by silica gel chromatography with hexane/ethyl acetate (1:4) and subsequent recrystallization from ethyl acetate/hexane gave 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene 35 as a colorless solid: mp 112.5-113.5°C; NMR (CDCl₃) δ 2.03-2.16 (m, 2H), 2.05 (s, 3H), 2.74-2.82 (m, 2H), 2.91-3.01 (m, 2H), 2.98 (s, 3H), 6.83-6.93 (m, 2H);

6.97-7.04 (m, 1H), 7.19 (d, $J = 8$ Hz, 2H), 7.68 (d, $J = 8$ Hz, 2H); MS (FAB) m/e (rel intensity) 337 (100), 331 (46); HRMS. Calc'd for $C_{19}H_{19}FO_2S$: 330.1090. Found: 330.1096. Anal. Calc'd for $C_{19}H_{19}FO_2S$: C, 69.07; H, 5.80; F, 5.75; S, 9.70. Found: C, 69.37; H, 5.81; F, 5.40; S, 9.78.

Example 3



10 1-[2-(4-Chlorophenyl)cyclopenten-1-yl]-
 4-(methylsulfonyl)benzene

Step 1: Preparation of 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene.

15

Following the synthetic procedure outlined in Step 3 of Example 1, 250 mg (0.93 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1, Step 2) was reacted with 300 mg (1.9 mmol) of 4-chlorophenylboronic acid (Lancaster). Purification by silica gel chromatography (MPLC) with hexane gave 290 mg of 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene as a colorless solid: mp 72-74°C; NMR ($CDCl_3$) δ 2.04 (m, $J = 7$ Hz, 2H), 2.46 (s, 3H), 2.86 (t, $J = 7$ Hz, 4H), 7.07-7.21 (m, 8H).


25

Step 2: Preparation of 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

30

Following the synthetic procedure outlined in Step 4 of Example 1, 280 mg (0.93 mmol) of 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene (Step 1) was oxidized. Purification by silica gel chromatography (MPLC) with hexane and subsequent recrystallization from ethyl acetate/hexane gave 192 mg (62%) of 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene as a colorless solid: mp 127.5-128.5°C; NMR (CDCl₃) δ 2.09 (m, J = 7 Hz, 2H), 2.91 (t, J = 7 Hz, 4H), 3.04 (s, 3H), 7.06, (d, J = 8 Hz, 2H), 7.21 (d, J = 8 Hz, 2H), 7.32 (d, J = 8 Hz, 2H), 7.77 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 332 (100), 297 (6), 218 (30); HRMS. Calc'd for C₁₈H₁₇ClO₂S: 332.0638. Found: 332.0628. Anal. Calc'd for C₁₈H₁₇ClO₂S: C, 64.95; H, 5.15; Cl, 10.65; S, 9.63. Found: C, 64.97; H, 5.15; Cl, 10.50; S, 9.58.

Example 4



1-[2-(2,4-Dichlorophenyl)cyclopenten-1-yl]-
4-(methylsulfonyl)benzene

Step 1: Preparation of 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene.

Following the synthetic procedure outlined in Step 3 of Example 1, 280 mg (0.93 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1, Step 2) was reacted with 350 mg (1.9 mmol) of 2,4-dichlorophenylboronic acid (Lancaster). Purification by silica gel chromatography (MPLC) with hexane gave 280 mg of 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene as an oil: NMR (CDCl₃) δ 2.10 (m, J = 7 Hz, 2H), 2.41 (s, 3H), 2.81 (t, J = 8 Hz, 4H), 2.92 (m, J = 8 Hz, 2H), 6.95-7.21 (m, 6H), 7.40 (s, 1H).


Step 2: Preparation of 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

15

Following the synthetic procedure outlined in Step 4 of Example 1, 280 mg (0.85 mmol) of 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylthio)benzene (Step 1) was oxidized. Purification by silica gel chromatography (MPLC) with hexane and subsequent lyophilization from acetonitrile/water (1:1) gave 158 mg (51%) of 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene as a solid: NMR (CDCl₃) δ 2.13 (m, J = 8 Hz, 2H), 2.84 (t, J = 8 Hz, 2H), 2.96 (t, J = 8 Hz, 2H), 3.00 (s, 3H), 6.97 (d, J = 8 Hz, 1H), 7.14-7.23 (m, 4H), 7.42 (d, J = 2 Hz, 1H), 7.71 (d, J = 9 Hz, 2H); MS (EI) m/e (rel intensity) 368 (61), 366 (91), 252 (100) 215 (64), 128 (47); HRMS. Calc'd for C₁₈H₁₆Cl₂O₂S: 366.0242. Found: 332.0249. Anal.

Calc'd for C₁₈H₁₆Cl₂O₂S: C, 58.86; H, 4.39; Cl, 19.37; S, 8.73. Found: C, 58.43; H, 4.47; Cl, 19.45; S, 8.82.

Example 5



1-[2-(4-Methylphenyl)cyclopenten-1-yl]- 4-(methylsulfonyl)benzene

5

Step 1: Preparation of 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene.

Following the synthetic procedure outlined in
 10 Step 3 of Example 1, 250 mg (0.93 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1, Step 2) was reacted with 260 mg (1.9 mmol) of 4-methylphenylboronic acid (Lancaster). Purification by silica gel chromatography (MPLC) with hexane gave 240 mg
 15 (92%) of 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene as a colorless solid: mp 64.5-66.5°C; NMR (CDCl₃) δ 2.03 (m, J = 7 Hz, 2H), 2.30 (s, 3H), 2.45 (s, 3H), 2.86 (t, J = 7 Hz, 4H), 6.99-7.15 (m, 8H).


20

Step 2: Preparation of 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

Following the synthetic procedure outlined in
 25 Step 4 of Example 1, 210 mg (0.75 mmol) of 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene (Step 1) was oxidized. Purification by silica gel chromatography (MPLC) with ethyl acetate/hexane (1:4) and subsequent recrystallization gave 140 mg (60%) of
 30 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-

(methylsulfonyl)benzene as a colorless solid: mp 118.0-118.5°C; NMR (CDCl₃) δ 2.07 (m, J = 7 Hz, 2H), 2.32 (s, 3H), 2.90 (t, J = 7 Hz, 4H), 3.03 (s, 3H), 6.99-7.08 (m, 4H), 7.34 (d, J = 8 Hz, 2H), 7.74 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 312 (100), 297 (10), 233 (11) 218 (22); HRMS. Calc'd for C₁₉H₂₀O₂S: 312.1184. Found: 312.1194. Anal. Calc'd for C₁₉H₂₀O₂S: C, 73.04; H, 6.45; S, 10.26. Found: C, 73.22; H, 6.65; S, 10.24.

Example 6



1-[2-(4-Trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

Step 1: Preparation of 4-trifluoromethylphenylboronic acid.

Following the synthetic procedure outlined in Step 1 of Example 1, 4-bromobenzotrifluoride (Aldrich) was converted to 4-trifluoromethyl phenylboronic acid: NMR (DMSO-d₆) δ 7.68 (d, J = 8 Hz, 2H), 7.98 (d, J = 8 Hz, 2H).

Step 2: Preparation of 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene.

Following the synthetic procedure outlined in Step 3 of Example 1, 240 mg (0.89 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1,

Step 2) was reacted with 360 mg (1.8 mmol) of 4-trifluoromethylphenylboronic acid (Step 1).

Purification by silica gel chromatography (MPLC) with hexane gave 240 mg (81%) of 1-[2-(4-

5 trifluoromethylphenyl)cyclopenten-1-yl]-4-

(methylthio)benzene as a colorless solid: mp 60.0-61.5°C; NMR (CDCl₃) δ 2.06 (m, J = 7 Hz, 2H), 2.46 (s, 3H), 2.89 (t, J = 7 Hz, 4H), 7.06 (d, J = 6 Hz, 2H), 7.10 (d, J = 6 Hz, 2H), 7.27 (d, J = 8 Hz, 2H), 7.46 (d, J = 8 Hz, 2H).

Step 3: Preparation of 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

15 Following the synthetic procedure outlined in Step 4 of Example 1, 215 mg (0.75 mmol) of 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-

(methylthio)benzene (Step 2) was oxidized. Purification by silica gel chromatography (MPLC) with ethyl


20 acetate/hexane (1:4) and subsequent recrystallization gave 163 mg (69%) of 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-

(methylsulfonyl)benzene as a colorless solid: mp 134.5-135.0°C; NMR (CDCl₃) δ 2.12 (m, J = 7 Hz, 2H), 2.95 (t,

25 J = 7 Hz, 4H), 3.05 (s, 3H), 7.24 (d, J = 8 Hz, 2H), 7.31 (d, J = 8 Hz, 2H), 7.49 (d, J = 8 Hz, 2H), 7.78 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 312 (100), 297 (10), 233 (11) 218 (22); HRMS. Calc'd for C₁₉H₂₀O₂S:

312.1184. Found: 312.1194. Anal. Calc'd for C₁₉H₂₀O₂S: C, 73.04; H, 6.45; S, 10.26. Found: C, 73.22; H, 6.65; S, 10.24.

Example 7



1-(2-Phenylcyclopenten-1-yl)-4-
(methylsulfonyl)benzene

5

Step 1: Preparation of 1-bromo-2-phenylcyclopentene.

Following a synthetic procedure which was similar to the one outlined in Step 2 of Example 1, 4.40 g (19.4 mmol) of 1,2-dibromocyclopentene was reacted with 2.0 g (17.7 mmol) of phenylboronic acid (Aldrich). Purification by silica gel chromatography (Waters Prep-500) with hexane gave 1.61 g (42%) of 1-bromo-2-phenylcyclopentene as an oil: NMR (CDCl₃) δ 2.01-2.10 (m, 2H), 2.74-2.82 (m, 2H), 2.82-2.90 (m, 2H), 7.27-7.33 (m, 1H), 7.33-7.41 (m, 2H), 7.57-7.63 (m, 2H).

15

Step 2: Preparation of 1-(2-(phenylcyclopenten-1-yl))-4-(methylthio)benzene.

20


Following a synthetic procedure which was similar to the one outlined in Step 3 of Example 1, 750 mg (3.4 mmol) of 1-bromo-2-phenylcyclopentene (Step 1) was reacted with 1.2 g (6.8 mmol) of 4-methylthiophenylboronic acid (Example 1, Step 1). Purification by silica gel chromatography (MPLC) with hexane gave 800 mg (89%) of 1-(2-phenylcyclopenten-1-yl)-4-(methylthio)benzene as an oil: NMR (CDCl₃) δ 2.00-2.17 (m, 2H), 2.46 (s, 3H), 2.86-3.01 (m, 4H), 7.08-7.19 (m, 4H), 7.19-7.32 (m, 5H).

30

Step 3: Preparation of 1-(2-(phenylcyclopenten-1-yl))-4-(methylsulfonyl)benzene.

Following the synthetic procedure outlined in Step 4 of Example 1, 800 mg (3.0 mmol) of 1-(2-phenylcyclopenten-1-yl)-4-(methylthio) benzene (Step 2) was oxidized. Purification by silica gel chromatography (MPLC) with hexane gave 300 mg (30%) of 1-(2-phenylcyclopenten-1-yl)-4-(methylsulfonyl)benzene as a colorless solid: mp 135.5-137.0°C; NMR (DMSO-d₆) δ 2.01 (m, J = 7 Hz, 2H), 2.91 (t, J = 7 Hz, 4H), 3.18 (s, 3H), 7.12-7.18 (m, 2H), 7.18-7.31 (m, 3H), 7.37 (d, J = 8 Hz, 2H), 7.76 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 298 (100), 219 (30), 141 (30); HRMS. Calc'd for C₁₈H₁₈O₂S: 298.1028. Found: 298.1056. Anal. Calc'd for C₁₈H₁₈O₂S: C, 72.45; H, 6.08; S, 10.74. Found: C, 72.46; H, 6.17; S, 10.56.

Example 8



1-[2-(4-Cyanophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

Step 1: Preparation of 4-cyanophenylboronic acid.

Following the synthetic procedure outlined in Step 1 of Example 1, 4-bromobenzonitrile (Aldrich) was converted to 4-cyanophenylboronic acid: NMR (DMSO-d₆) δ 7.76 (d, J = 7 Hz, 2H), 7.91 (d, J = 8 Hz, 2H).


Step 2: Preparation of 1-[2-(4-cyanophenyl)
cyclopenten-1-yl]-4-(methylthio)benzene.

Following the synthetic procedure outlined in
5 Step 3 of Example 1, 500 mg (1.9 mmol) of 1-(2-
bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1,
Step 2) was reacted with 540 mg (3.7 mmol) of 4-
cyanophenylboronic acid (Step 1). Purification by
silica gel chromatography (MPLC) with hexane/ethyl
10 acetate (19:1) gave 480 mg (89%) of 1-[2-(4-
cyanophenyl)cyclopenten-1-yl]-4-(methylthio)benzene as
an oil: NMR (CDCl₃) δ 2.07 (m, J = 7 Hz, 2H), 2.42 (s,
3H), 2.89 (t, J = 7 Hz, 4H), 7.05 (d, J = 8 Hz, 2H), 7.11
(d, J = 8 Hz, 2H), 7.26 (d, J = 8 Hz, 2H), 7.48 (d, J = 8
15 Hz, 2H).

Step 3: Preparation of 1-[2-(4-cyanophenyl)
cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

20 Following the synthetic procedure outlined in
Step 4 of Example 1, 240 mg (0.82 mmol) of 1-[2-(4-
cyanophenyl)cyclopenten-1-yl]-4-(methylthio)benzene
(Step 2) was oxidized. Purification by silica gel
chromatography (MPLC) with ethyl acetate/hexane (3:7)
25 and subsequent recrystallization gave 174 mg (66%) of 1-
[2-(4-cyanophenyl) cyclopenten-1-yl]-4-
(methylsulfonyl)benzene as a colorless solid: mp 163.0-
164.5°C; NMR (CDCl₃) δ 2.13 (m, J = 7 Hz, 2H), 2.95 (t,
 J = 7 Hz, 4H), 3.05 (s, 3H), 7.22 (d, J = 8 Hz, 2H), 7.30
30 (d, J = 8 Hz, 2H), 7.51 (d, J = 8 Hz, 2H), 7.79 (d, J = 8
Hz, 2H); MS (EI) m/e (rel intensity) 323 (100), 308 (4),
244 (42); HRMS. Calc'd for C₁₉H₁₇NO₂S: 323.0980.
Found: 323.1014. Anal. Calc'd for C₁₉H₁₇NO₂S: C,
70.56; H, 5.30; N, 4.33; S, 9.91. Found: C, 70.59; H,
35 5.34; N, 4.29; S, 9.67.

Example 9



1-[2-(4-Hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

5

Step 1: Preparation of 4-hydroxymethylphenylboronic acid.

Following the synthetic procedure outlined in Step 1 of Example 1, 4-bromobenzyl alcohol (Aldrich) was converted to 4-hydroxymethylphenyl boronic acid: NMR (DMSO- d_6) δ 4.50 (d, J = 5 Hz, 2H), 5.08 (s, 1H), 7.20-7.45 (m, 2H), 7.68-7.90 (m, 2H).

Step 2: Preparation of 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene.


Following the synthetic procedure outlined in Step 3 of Example 1, 250 mg (0.93 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1, Step 2) was reacted with 290 mg (1.9 mmol) of 4-hydroxymethylphenylboronic acid (Step 1). Purification by silica gel chromatography (MPLC) with ethyl acetate/hexane (1:4) gave 255 mg (92%) of 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene as a solid: mp 82-85°C; NMR (CDCl₃) δ 2.04 (m, J = 7 Hz, 2H), 2.45 (s, 3H), 2.88 (t, J = 7 Hz, 4H), 4.64 (s, 2H), 7.08 (s, 4H), (m, 2H), 7.15-7.25 (m, 4H).

30

Step 3: Preparation of 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

Following the synthetic procedure outlined in Step 4 of Example 1, 210 mg (0.71 mmol) of 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylthio)benzene (Step 2) was oxidized. Purification by silica gel chromatography with hexane/ethyl acetate (1:4) and subsequent recrystallization from ethyl acetate/hexane gave 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene as a colorless solid: mp 114-115°C; NMR (CDCl₃) δ 2.09 (m, J = 7 Hz, 2H), 2.92 (t, J = 7 Hz, 4H), 3.03 (s, 3H), 4.67 (s, 2H), 7.13 (d, J = 8 Hz, 2H), 7.24 (d, J = 8 Hz, 1H), 7.33 (d, J = 8 Hz, 2H), 7.74 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 328 (100), 311 (10), 297 (21), 218 (53); HRMS. Calc'd for C₁₉H₂₀O₃S: 328.1133. Found: 328.1147. Anal. Calc'd for C₁₉H₂₀O₃S: C, 69.48; H, 6.14; S, 9.76. Found: C, 69.51; H, 6.40; S, 9.68.

Example 10




1-[2-(4-Methoxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

Under nitrogen, a stirred solution of 79 mg (0.24 mmol) of 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (Example 9) in 2 mL of dry THF at 0°C was treated with 15 mg (0.6 mmol) of sodium hydride (95%). After 30 minutes, 0.1 mL (1.6

mmol) of methyl iodide was added and the reaction was allowed to warm to ambient temperature overnight. The solvent was removed in vacuo; the residue was dissolved in ethyl acetate and washed with water, dried (MgSO₄), and reconcentrated. Purification by silica gel chromatography (MPLC) with hexane/ethyl acetate (5:1) and subsequent lyophilization from acetonitrile/water (1:1) gave 25 mg (30%) of 1-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene as a colorless solid: NMR (CDCl₃) δ 2.09 (m, J = 7 Hz, 2H), 2.92 (t, J = 7 Hz, 4H), 3.03 (s, 3H), 3.40 (s, 3H), 4.42 (s, 2H), 7.12 (d, J = 8 Hz, 2H), 7.20 (d, J = 8 Hz, 2H), 7.33 (d, J = 8 Hz, 2H), 7.73 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 342 (100), 81 (27), 69 (62). HRMS. Calc'd for C₂₀H₂₂O₃S: 342.1290. Found: 342.1301. Anal. Calc'd for C₂₀H₂₂O₃S: C, 70.15; H, 6.48; S, 9.36. Found: C, 69.86; H, 6.64; S, 9.38.

Example 11



1-[2-(4-Methylthiophenyl)cyclopenten-1-yl]- 4-(methylsulfonyl)benzene

Step 1: Preparation of 1-(2-bromocyclopenten-1-yl)-4-(methylsulfonyl)benzene.

Following the synthetic procedure outlined in Step 4 of Example 1, 250 mg (0.93 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylthio)benzene (Example 1, Step 2) was oxidized to give 280 mg (100%) of 1-(2-


bromocyclopenten-1-yl)-4-(methylsulfonyl)benzene as a colorless solid: mp 103-104°C; NMR (CDCl₃) δ 2.02-2.14 (m, 2H), 2.74-2.83 (m, 2H), 2.86-2.94 (m, 2H), 3.07 (s, 3H), 7.77 (d, *J* = 8 Hz, 2H), 7.93 (d, *J* = 8 Hz, 2H).

5

Step 2: Preparation of 1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene.

Following a synthetic procedure which was similar to the one outlined in Step 3 of Example 1, 270 mg (0.9 mmol) of 1-(2-bromocyclopenten-1-yl)-4-(methylsulfonyl)benzene (Step 1) was reacted with 300 mg (1.8 mmol) of 4-methylthiophenylboronic acid (Example 1, Step 1). Purification by silica gel chromatography (MPLC) with hexane/ethyl acetate (4:1) gave 265 mg (86%) of 1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene as a colorless solid: mp 138-139°C; NMR (CDCl₃) δ 2.08 (m, *J* = 7 Hz, 2H), 2.47 (s, 3H), 2.90 (t, *J* = 7 Hz, 4H), 3.04 (s, 3H), 7.06 (d, *J* = 8 Hz, 2H), 7.11 (d, *J* = 8 Hz, 2H), 7.34 (d, *J* = 8 Hz, 2H), 7.75 (d, *J* = 8 Hz, 2H); MS (EI) *m/e* (rel intensity) 344 (100), 297 (4), 218 (33); HRMS. Calc'd for C₁₉H₂₀O₂S₂: 344.0905. Found: 344.0907. Anal. Calc'd for C₁₉H₂₀O₂S₂: C, 56.24; H, 5.85; S, 18.16. Found: C, 56.28; H, 5.81; S, 18.95.

Example 12



1-[2-(4-Fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene

30

Step 1: Preparation of the ethyl acetal of mucobromic acid.

5 Under nitrogen, a stirred solution of 500 g (1.94 mol) of mucobromic acid (Lancaster) and 2 g of p-toluenesulfonic acid monohydrate in 600 mL of toluene and 400 mL of absolute ethanol was heated to reflux for 6 hours during which time 150 mL of a water, toluene, and ethanol azeotrope was removed by distillation. The solution was concentrated in vacuo; the residue was dissolved in 1500 mL of ethyl acetate and washed successively with water, saturated sodium carbonate, and brine, dried (Na₂SO₄), and reconcentrated to give 440 g (79%) of the ethyl acetal of mucobromic acid (5 in Synthetic Scheme II) as an oil: NMR (CDCl₃) δ 1.31 (t, J= 7 Hz, 3H), 3.73-3.96 (m, 2H), 5.81 (s, 1H).

20 Step 2: Preparation of cis-2,3-dibromobut-2-ene-1,4-diol.

Under nitrogen, a stirred solution of 150 g (525 mmol) of the ethyl acetal of mucobromic acid (Step 1) in 150 mL of anhydrous THF at -78°C was treated with 1400 mL of diisobutylaluminum hydride (1.5 M in toluene) over a 30 minute period. The solution was allowed to warm to ambient temperature and stirred for 2 hours. The reaction was slowly treated (maintaining the temperature below 10°C) with 100 mL of acetone followed by 50 mL of 2.5 N sodium hydroxide. Water (1000 mL) was added and the solution extracted 5 times with ethyl acetate. The combined extracts were washed with brine, dried (Na₂SO₄), and concentrated. The residue was washed 3 times with hexane and dried in vacuo to give 88.5 g (69%) of cis-2,3-dibromo-2-ene-1,4-diol (6 in Synthetic Scheme II) as a colorless solid: mp 66-67°C; NMR (DMSO-d₆) δ 4.27 (d, J=6 Hz, 4H), 5.44 (t, J= 6 Hz, 2H).

Step 3: Preparation of cis-1,2,3,4-tetrabromobut-2-ene

Under nitrogen, a stirred solution of 25.2 g
5 (102 mmol) of cis-2,3-dibromobut-2-ene-1,4-diol (Step 2)
in 150 mL of methylene chloride at 0°C was treated with
9.6 mL of phosphorus tribromide. The solution was
allowed to warm to ambient temperature where it was
allowed to stir for 1 hour prior to the addition of ice
10 water. The aqueous phase was extracted 5 times with
methylene chloride; these extracts were combined with
the original methylene chloride phase and washed with
water, saturated sodium carbonate, brine, dried
(Na₂SO₄), and concentrated in vacuo to give 18 g (47%)
15 of cis-1,2,3,4-tetrabromobut-2-ene [7 (X= Br) in
Synthetic Scheme II] as an oil: NMR (CDCl₃) δ 4.40 (s,
4H).

Step 4: Preparation of 1,2-dibromo-4,4-di(carboethoxy)
20 cyclopentene.

Under nitrogen, a solution of 9.7 g (60.6
mmol) of diethyl malonate in anhydrous THF at -10°C was
treated with 2.9 g (121 mmol) of sodium hydride (95%)
25 and allowed to stir for 30 minutes. The resulting
solution was then added slowly to 15 g (40.4 mmol) of
cis-1,2,3,4-tetrabromobut-2-ene (Step 3) in 350 mL of
anhydrous THF at -78°C. The reaction was allowed to
warm to ambient temperature overnight prior to
30 concentration in vacuo. The residue was dissolved in
ethyl acetate and washed with water, dried (Na₂SO₄), and
reconcentrated. Purification by silica gel
chromatography (Waters Prep-500) with ethyl
acetate/hexane (1:99) gave 3.7 g (25%) of 1,2,-dibromo-
35 4,4-di(carboethoxy)cyclopentene (8 in Synthetic Scheme
II) as a colorless oil: NMR (CDCl₃) δ 1.26 (t, J= 7 Hz,
6H), 3.26 (s, 4H), 4.22 (m, J= 7 Hz, 4H); MS (FAB) for
M+H m/e: 373, 371, 369.

Step 5: Preparation of 1,2-dibromo-4,4-di(hydroxymethyl)cyclopentene.

5 Under nitrogen, a stirred solution of 8.7 g (23.5 mmol) of 1,2-dibromo-4,4-di(carboethoxy)cyclopentene (Step 4) in 70 mL of anhydrous THF at -78°C was treated with 80 mL of diisobutylaluminum hydride (1.5 M in toluene) over a 20
10 minute period. The reaction was allowed to warm to ambient temperature overnight and was slowly treated with 20 mL of acetone followed by 10 mL of 2.5 N sodium hydroxide. Water (100 mL) was added and the solution extracted 5 times with ethyl acetate. The combined
15 extracts were washed with brine, dried (Na₂SO₄), and concentrated in vacuo to give 5.7 g (85%) of 1,2-dibromo-4,4-di(hydroxymethyl)cyclopentene (10 in Synthetic Scheme III) as a colorless oil: NMR (CDCl₃) δ 2.20 (s, 2H), 2.50 (s, 4H), 3.70 (s, 4H).

20

Step 6: Preparation of 1,2-dibromo-4,4-di(tosylmethyl)cyclopentene.

Under nitrogen, a stirred solution of 5.7 g (19.9 mmol) of 1,2-dibromo-4,4-di(hydroxymethyl)cyclopentene (Step 5) in 50 mL of pyridine at ambient temperature was treated with 19 g (99.7 mmol) of p-toluenesulfonyl chloride. The reaction was allowed to stir overnight and was concentrated in vacuo. The residue was dissolved ethyl acetate and
30 washed twice with 3% hydrochloric acid followed by brine. The solution was dried (Na₂SO₄) and concentrated in vacuo to give 5.2 g (44%) of 1,2-dibromo-4,4-di(tosylmethyl)cyclopentene (14 in Synthetic Scheme V) as a colorless semisolid: NMR (CDCl₃) δ 2.42 (s, 4H), 2.47 (s, 6H), 3.90 (s, 4H), 7.37 (d, J = 8 Hz, 4H), 7.74 (d, J = 8 Hz, 4H).

35

Step 7: Preparation of 1,2-dibromo-4,4-di(iodomethyl)cyclopentene.

Under nitrogen, a stirred solution of 5.2 g (8.7 mmol) of 1,2-dibromo-4,4-di(tosylmethyl)cyclopentene (Step 6) and 13 g (86 mmol) of sodium iodide in 40 mL of DMF/H₂O (3:1) was heated to 150°C in an oil bath overnight. The reaction was cooled, diluted with 200 mL of ethyl acetate, and washed with water. Drying (Na₂SO₄) and concentrating in vacuo gave 3.7 g (84%) of 1,2-dibromo-4,4-di(iodomethyl)cyclopentene (15 in Synthetic Scheme V) as an oil: NMR (CDCl₃) δ 2.70 (s, 4H), 3.50 (s, 4H).

Step 8: Preparation of 1,2-dibromo-4,4-dimethylcyclopentene.

Under nitrogen, a stirred solution of 3.7 g (7.3 mmol) of 1,2-dibromo-4,4-di(iodomethyl)cyclopentene (Step 7) and 1.3 g (20.6 mmol) of sodium cyanoborohydride in 15 mL of hexamethylphosphoramide (HMPA) was heated to 100 °C in an oil bath overnight. The reaction was cooled, diluted with 50 mL of water, and extracted 5 times with ethyl acetate/hexane (1:5). The combined extracts were washed 3 times with water, dried (Na₂SO₄), and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500) with hexane gave 1.3 g (70%) of 1,2-dibromo-4,4-dimethylcyclopentene (16 in Synthetic Scheme V) as a colorless oil: NMR (CDCl₃) δ 1.16 (s, 6H), 2.44 (s, 4H); MS (EI) m/e (rel intensity) 256 (24), 254 (63), 252(44), 175 (26), 173 (29), 94 (100).

Step 9: Preparation of 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylthio)benzene.


Under nitrogen, 1.3 g (5.1 mmol) of 1,2-dibromo-4,4-dimethylcyclopentene (Step 8) was reacted with 600 mg (4.3 mmol) of 4-fluorophenylboronic acid (Lancaster) in 23 mL of toluene, 15 mL of ethanol, and 10 mL of 2M Na₂CO₃ in the presence of 250 mg (5 mol %) of Pd(PPh₃)₄. The reaction was vigorously stirred at reflux overnight and concentrated in vacuo. The residue was dissolved in ethyl acetate and washed with water, dried (Na₂SO₄), and reconcentrated. Purification by silica gel chromatography (MPLC) with hexane gave 250 mg of 1-(2-bromocyclopenten-1-yl)-4-fluorobenzene (38 in Synthetic Scheme X when R¹ and R² = CH₃, R⁵ = F, R³, R⁴, R⁶, and R⁷ = H) as a pale yellow oil which was reacted with 200 mg (1.2 mmol) of 4-methylthiophenylboronic acid (Example 1, Step 1) in 5.2 mL of toluene, 3.4 mL of ethanol, and 2.2 mL of 2M Na₂CO₃ in the presence of 40 mg (5 mol %) of Pd(PPh₃)₄. The reaction was vigorously stirred at reflux for 6 hours and concentrated in vacuo. The residue was dissolved in ethyl acetate and washed with water, dried (Na₂SO₄), and reconcentrated. Purification by silica gel chromatography (MPLC) with hexane gave 120 mg of 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylthio)benzene (41 in Synthetic Scheme X when R¹ and R²=CH₃, R⁵= F, R¹⁰= SCH₃, R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹¹, and R¹²= H) as an oil: NMR (CDCl₃) δ 1.20 (s, 6H), 2.42 (s, 3H), 2.63 (s, 4H), 6.90 (t, J= 8 Hz, 2H), 7.00-7.18 (m, 4H), 7.30-7.60 (m, 2H).

Step 10: Preparation of 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene.

A solution of 120 mg (0.39 mmol) of 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylthio)benzene (Step 9) in 3 mL of methanol/water (1:1) was slowly treated with 470 mg (0.76 mmol) of Oxone® [2 KHSO₅.KHSO₄.K₂SO₄] in 2 mL of water. After stirring for 4 hours, the solvent was removed in vacuo.

The residue was dissolved in ethyl acetate and washed with water and brine, dried (MgSO₄), and reconcentrated. Purification by silica gel chromatography (MPLC) with hexane/ethyl acetate (5:1) and subsequent lyophilization from acetonitrile/water (1:1) gave 50 mg (38%) of 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene (41 in Synthetic Scheme X when R¹ and R²= CH₃, R⁵= F, R¹⁰= SO₂CH₃, R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹¹, and R¹²= H) as a colorless solid: NMR (CDCl₃) d 1.24 (s, 6H), 2.71 (s, 4H), 3.4 (s, 3H), 6.92 (t, *J*= 8 Hz, 2H), 7.05-7.12 (m, 2H), 7.30 (d, *J*= 9 Hz, 2H), 7.75 (d, *J*= 9 Hz, 3H); MS (EI) m/e (rel intensity) 344 (100), 329 (33), 250 (18), 235 (20), 109 (35), 69 (44); HRMS. Calc'd for C₂₀H₂₁FO₂S: 344.1246. Found: 344.1272.

Example 13




4-[2-(4-Fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide

20

Under nitrogen, a solution of 4.55 g (13.2 mmol) of 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 12) in 50 mL of THF at -78°C was treated with 25 6.3 mL (15.8 mmol) of *n*-butyllithium (2.5 M in hexane) over a period of 5 minutes. The reaction was stirred at ambient temperature for 25 minutes, cooled to -78°C, and treated with 19.8 mL (19.8 mmol) of tributylborane (1.0 M in THF). The resulting dark brown solution was 30 stirred at ambient temperature for 20 minutes and then

at reflux for 16 hours prior to the addition of 8.7 g (106 mmol) of sodium acetate, 10 mL of water, and 5.2 g (46 mmol) of hydroxylamine-O-sulfonic acid. The resulting light orange mixture was stirred at ambient temperature for 3 hours and the aqueous phase extracted with ethyl acetate. The combined extracts were washed first with water and then with brine, dried (MgSO₄), and concentrated in vacuo. The residue was chromatographed on silica gel to give 2.0 g (44%) of 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 117-118°C; NMR (CDCl₃) δ 1.23 (s, 6H), 2.70 (s, 4H), 4.72 (s, 2H), 6.92 (t, J = 9 Hz, 2H), 7.05-7.12 (m, 2H), 7.26 (d, J = 9 Hz, 2H), 7.74 (d, J = 9 Hz, 2H); MS (EI) m/e (rel intensity) 345 (100), 330 (39), 250 (28), 109 (50), 80 (31), 69 (58); HRMS. Calc'd for C₁₉H₂₀FNO₂S: 345.1199. Found: 345.1194. Anal. Calc'd for [C₁₉H₂₀FNO₂S + 0.27 H₂O]: C, 65.15; H, 5.91; N, 4.00; S, 9.15. Found: C, 65.07; H, 5.94; N, 3.86; S, 9.29.

Example 14



1-[2-(4-Chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene

Following the general procedures outlined in Synthetic Schemes XV, XX, XXI, and XXII, 1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene was isolated as a colorless solid: mp 144-145°C; NMR (CDCl₃) δ 1.23 (s, 6H), 2.71


106

(s, 4H), 3.04 (s, 3H), 7.05 (d, J = 8 Hz, 2H), 7.21 (d, J = 8 Hz, 2H), 7.30 (d, J = 8 Hz, 2H), 7.76 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 360 (100), 345 (30), 266 (20), 231 (20); HRMS. Calc'd for $C_{20}H_{21}ClLiO_2S$:

- 5 367.1111. Found: 367.1105. Anal. Calc'd for $C_{20}H_{21}ClO_2S$: C, 66.50; H, 5.82; Cl, 9.84. Found: C, 66.61; H, 6.10, Cl, 9.85.

Example 15

10




4-[2-(4-Chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide

15

- Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl) benzene (the title compound of Example 14) was converted to 4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 144-145°C; NMR ($CDCl_3$) δ 1.24 (s, 6H), 2.71 (s, 4H), 4.78 (s, 2H), 7.05 (d, J = 9 Hz, 2H), 7.20 (d, J = 9 Hz, 2H), 7.26 (d, J = 9 Hz, 2H), 7.75 (d, J = 9 Hz, 2H); HRMS. Calc'd for $C_{19}H_{20}ClNO_2S$: 361.0903. Found: 361.0882. Anal. Calc'd for $C_{19}H_{20}ClNO_2S$: C, 62.72; H, 5.55; Cl, 10.31; S, 8.80. Found: C, 62.35; H, 5.71; Cl, 10.04; S, 8.74.

30

Example 16



5

1-[2-(4-Fluorophenyl)-4,4-
di(fluoromethyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene

10 Step 1: Preparation of 4-(methylthio)acetophenone

To a stirred solution of 50 g (340 mmol) of 4-(methylthio)benzonitrile in 2 L of THF at -78°C under an atmosphere of nitrogen, was added 282 mL (390 mmol) of methyl lithium (1.4 M in diethyl ether) over a period of ten minutes. The solution was stirred at -78°C for one hour, and then the dry ice bath was removed. After five hours, 100 mL of water followed by 200 mL of 3N hydrochloric acid were added to the reaction mixture and it was stirred overnight. Concentration in vacuo gave a residue which was partitioned between ethyl acetate and water. The water layer was extracted with three portions of ethyl acetate and the combined ethyl acetate layers were dried (MgSO_4). Concentration in vacuo gave 58 g of crude 4-(methylthio)acetophenone as a solid: NMR (CDCl_3) δ 2.52 (s, 3H), 2.57 (s, 3H), 7.26 (d, $J = 9$ Hz, 2H), 7.87 (d, $J = 9$ Hz, 2H).

30 Step 2: Preparation of 4-(methylsulfonyl)acetophenone

To a solution of 11.73 g (71.1 mmol) of 4-(methylthio)acetophenone (prepared in Step 1) in 500 mL

of dichloromethane at ambient temperature was added 61.14 g (177 mmol) of *m*-chloroperoxybenzoic acid (50%) (MCPBA) in portions over 20 minutes. The reaction was stirred for two hours, quenched slowly with aqueous sodium bisulfite, washed with three 100 mL portions of saturated sodium bicarbonate, dried (MgSO_4), and concentrated in vacuo to give 11.91 g (91%) of 4-(methylsulfonyl)acetophenone as a colorless solid: NMR (CDCl_3) δ 2.67 (s, 3H), 3.08 (s, 3H), 8.06 (d, J = 9 Hz, 2H), 8.14 (d, J = 9 Hz, 2H).

Step 3: Preparation of 2-bromo-4'-(methylsulfonyl)acetophenone

To a stirred solution of 11.91 g (60.5 mmol) of 4-(methylsulfonyl)acetophenone (prepared in Step 2) in 133 mL of glacial acetic acid and 0.11 mL of hydrochloric acid at ambient temperature was added a solution of 8.22 g (51.4 mmol) of bromine in 9.3 mL of glacial acetic acid over a period of three hours. The reaction mixture was diluted with 500 mL of water and extracted with chloroform. The combined extracts were dried (MgSO_4) and concentrated in vacuo to give 15.7 g of crude 2-bromo-4'-(methylsulfonyl)acetophenone as a solid: NMR (CDCl_3) δ 3.10 (s, 3H), 4.45 (s, 2H), 8.08 (d, J = 9 Hz, 2H), 8.17 (d, J = 9 Hz, 2H).

Step 4: Preparation of 2-(4-fluorophenyl)-1-[2-[4-(methylsulfonyl)phenyl]-2-oxoethoxy]ethanone

To a stirred solution of 4.45 g (28.9 mmol) of 4-fluorophenylacetic acid in 3.26 g (31.8 mmol) of triethylamine and 275 mL of acetonitrile was added 8.9 g (28.9 mmol) of 2-bromo-4'-(methylsulfonyl)acetophenone (prepared in Step 3) at ambient temperature. The reaction mixture was stirred for 30 minutes, concentrated in vacuo, and partitioned between ethyl

acetate and water. The organic phase was dried (MgSO_4) and concentrated in vacuo. Purification by silica gel chromatography with ethyl acetate/hexane (1:1) gave 6.87 g (68%) of 2-(4-fluorophenyl)-1-[2-[4-

- 5 (methylsulfonyl)phenyl]-2-oxoethoxy]ethanone as a colorless solid: NMR (CDCl_3) δ 3.08 (s, 3H), 3.79 (s, 2H), 5.35 (s, 2H), 7.06 (s, t, J = 9 Hz, 2H), 7.32 (dd, J = 6 and 9 Hz, 2H), 8.06 (s, 4H).

10 Step 5: Preparation of 3-(4-fluorophenyl)-4-[(4-methylsulfonyl)phenyl]-5H-furan-2-one

- Under nitrogen, 4.10 g (11.7 mmol) of 2-(4-fluorophenyl)-1-[2-[4-(methylsulfonyl)phenyl]-2-oxoethoxy]ethanone (prepared in Step 4), 6.52 mL (46.8 mmol) of triethylamine, 4.89 g (25.7 mmol) of *p*-toluenesulfonic acid, and 12 g of 4Å molecular sieves were added to 117 mL of acetonitrile and stirred at reflux for 16 hours. The reaction mixture was concentrated in vacuo and the residue partitioned between dichloromethane and water. The dichloromethane layer was dried (MgSO_4) and reconcentrated in vacuo.

- 20 Recrystallization from hexane/ethyl acetate (2:1) gave 3.65 g (94%) of 3-(4-fluorophenyl)-4-[(4-methylsulfonyl)phenyl]-5H-furan-2-one as a solid: mp 166-167°C; NMR (CDCl_3) δ 3.08 (s, 3H), 5.19 (s, 2H), 7.10 (t, J = 9 Hz, 2H), 7.42 (dd, J = 6 and 9 Hz, 2H), 7.52 (d, J = 9 Hz, 2H), 7.97 (d, J = 9 Hz, 2H); HRMS. Calc'd for $\text{C}_{17}\text{H}_{13}\text{FO}_4\text{S}$: 332.0519. Found: 332.0501.
- 30 Anal. Calc'd for $\text{C}_{17}\text{H}_{13}\text{FO}_4\text{S}$: C, 61.44; H, 3.94; O, 19.26. Found: C, 61.11; H, 4.06; O, 19.32.

Step 6: Preparation of 2-(4-fluorophenyl)-3-[(4-methylsulfonyl)phenyl]-1,4-dihydroxy-2-butene

35

To a solution of 3.08 g (9.28 mmol) of 3-(4-fluorophenyl)-4-(4-methylsulfonyl)phenyl]-5H-furan-2-one

110

(prepared in Step 5) in 93 mL of tetrahydrofuran (THF) at -78°C under an atmosphere of nitrogen was added 20 mL (30 mmol) of diisobutylaluminum hydride (1.5 M in THF) (DIBAL) over a 10 minute period. The solution was stirred at -78°C for 20 minutes, allowed to warm to ambient temperature, and stirred overnight. An additional 15 mL (22 mmol) aliquot of DIBAL was added and stirring was continued for 2 hours. The reaction was cooled to -78°C, treated dropwise with 25 mL of acetone, warmed to room temperature, and slowly treated with 25 mL of water. The mixture was stirred for 30 minutes prior to the careful addition of 35 mL of 1.2N sodium hydroxide. The mixture was extracted with ethyl acetate, washed with 1 N hydrochloric acid followed by brine, dried (MgSO₄), and concentrated in vacuo to give 3.8 g of crude 2-(4-fluorophenyl)-3-[(4-methylsulfonyl)phenyl]-1,4-dihydroxy-2-butene as a colorless oil: NMR (CDCl₃) δ 2.98 (s, 3H), 4.60 (d, J = 6 Hz, 4H), 6.8 (t, J = 9 Hz, 2H), 6.94-7.02 (m, 2H), 7.22 (d, J = 9 Hz, 2H), 7.65 (d, J = 9 Hz, 2H).

Step 7: Preparation of 2-(4-fluorophenyl)-3-[(4-methylsulfonyl)phenyl]-1,4-dichloro-2-butene

To a solution of 3.5 g (7.62 mmol) of crude 2-(4-fluorophenyl)-3-[(4-methylsulfonyl)phenyl]-1,4-dihydroxy-2-butene (prepared in Step 6) in 58 mL of N,N-dimethylformamide (DMF) at 5°C under an atmosphere of nitrogen was added dropwise 1.52 mL (20.84 mmol) of thionyl chloride. The reaction was stirred at 5°C for 22 hours, stirred at ambient temperature for an additional 8 hours, and concentrated in vacuo. The residue was partitioned between ethyl acetate and water; the ethyl acetate phase was dried (MgSO₄) and concentrated in vacuo to give crude 2-(4-fluorophenyl)-3-[(4-methylsulfonyl)phenyl]-1,4-dichloro-2-butene as a solid: NMR (CDCl₃) δ 3.0 (s, 3H), 4.55 (d, J = 3.4 Hz, 4H),

111

6.86 (t, $J = 9$ Hz, 2H), 6.75 (d, $J = 8.3$ Hz, 2H), 7.45 (d, $J = 9$ Hz, 2H).

5 Step 8A: Preparation of 1-[2-(4-fluorophenyl)-4,4-
 dicarbomethoxycyclopenten-1-yl]-4-
 (methylsulfonyl)benzene

To a solution of 1.2 mL (10.5 mmol) of dimethyl malonate in 10 mL of DMF under an atmosphere of
10 nitrogen was added 215 mg (26.9 mmol) of lithium hydride in portions. The resulting suspension was stirred at ambient temperature for 20 minutes prior to the addition of a solution of crude 2-(4-fluorophenyl)-3-[(4-methylsulfonyl)phenyl]-1,4-dichloro-2-butene (prepared
15 in Step 7) in 10 mL of DMF. The reaction was stirred at ambient temperature for 15 hours, treated with another 150 mg (18.8 mmol) of lithium hydride, and stirred for another 4 hours. The mixture was concentrated in vacuo and partitioned between ethyl acetate and water; the
20 organic phase was dried ($MgSO_4$), and concentrated in vacuo. The residue was chromatographed on silica gel to give 1.1 g (34%) of 1-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-4-(methylsulfonyl)benzene as an oil: NMR ($CDCl_3$) δ 3.03
25 (s, 3H), 3.55 (s, 4H), 3.79 (s, 6H), 6.93 (t, $J = 9$ Hz, 2H), 7.11 (dd, $J = 6$ and 9 Hz, 2H), 7.32 (d, $J = 9$ Hz, 2H), 7.77 (d, $J = 9$ Hz, 2H).

30 Step 8B: Preparation of 1-[2-(4-fluorophenyl)-4,4-
 dicarbomethoxycyclopenten-1-yl]-4-
 (methylsulfonyl)benzene

To a solution of 7.18 mL (63 mmol) of dimethyl malonate in 160 mL of DMF at 0°C under an atmosphere of
35 nitrogen was added 3.0 g (75 mmol) of sodium hydride (60% suspension in oil). The reaction was stirred at ambient temperature for 15 minutes (or until the gas

evolution has ceased), cooled to -20°C , and treated with 15 g (69 mmol) of 2-bromo-4'-fluoroacetophenone (Aldrich) in one portion. The mixture was stirred at ambient temperature for 1 hour and then cooled to 0°C ;
5 another 75 mmol of sodium hydride was added and the resulting mixture stirred at ambient temperature for 15 minutes (or until the gas evolution has ceased). The reaction was recooled to -20°C and treated with 19.1 g (69 mmol) of 2-bromo-4'-(methylsulfonyl)acetophenone
10 (prepared in Step 3). The reaction was stirred at room temperature for 2 hours and concentrated in vacuo. The residue was partitioned between water and ethyl acetate; the ethyl acetate phase was dried (MgSO_4) and reconcentrated in vacuo. The residue was chromatographed
15 on silica gel to give 13.8 g (51%) of dimethyl 2-[2-(4-fluorophenyl)-2-oxoethyl]-2-[2-[4-(methylsulfonyl)phenyl]-2-oxoethyl]propanedioate as an oil: NMR (CDCl_3) δ 3.06 (s, 3H), 3.76 (s, 6H), 4.03 (s, 2H), 4.08 (s, 2H), 7.13 (t, $J = 8.6$ Hz, 2H), 7.97-8.05
20 [m with d at 8.03 ($J = 8.7$ Hz), 4H], 8.14 (d, $J = 8.5$ Hz, 2H).

To a vigorously stirred mixture of 50.4 g (771 mmol) of zinc dust in 640 mL of THF at -78°C under an
25 atmosphere of nitrogen was added dropwise 60.4 mL (551 mmol) of titanium (IV) chloride. The reaction was warmed to ambient temperature with a water bath and then stirred at reflux for 1 hour. To the resulting dark mixture under reflux was added a solution of 15 g (32.3
30 mmol) of dimethyl 2-[2-(4-fluorophenyl)-2-oxoethyl]-2-[2-[4-(methylsulfonyl)phenyl]-2-oxoethyl]propanedioate (prepared above) in 20 mL of THF. The resulting mixture was stirred at ambient temperature for 16 hours, filtered through a pad of celite, rinsed with ethyl
35 acetate, and concentrated in vacuo. The residue was partitioned between water and ethyl acetate; the organic phase was washed with brine, dried (MgSO_4), and concentrated in vacuo. The residue was chromatographed

on silica gel to give 6.26 g (44%) of 1-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-4-(methylsulfonyl)benzene which was identical to the material prepared in Step 8, Method A.

5

Step 9: Preparation of 1-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

10

Under nitrogen, a solution of 1.01 g (2.34 mmol) of 1-[2-(4-fluorophenyl)-4,4-dicarbomethoxycyclopenten-1-yl]-4-(methylsulfonyl)benzene (prepared in Step 8) in 1.5 mL of THF at -78°C was treated with 11.6 mL (11.6 mmol) of DIBAL (1.0 M in THF). The reaction was stirred at ambient temperature for 1.5 hours, quenched with acetone and aqueous NaOH, extracted with ethyl acetate, dried (MgSO₄), and concentrated in vacuo to give 840 mg of crude 1-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene as a colorless oil: NMR (CDCl₃) δ 2.82 (d, J = 5 Hz, 4H), 3.04 (s, 3H), 3.86 (d, J = 5 Hz, 4H), 6.94 (t, J = 9 Hz, 2H), 7.11 (dd, J = 5 and 9 Hz, 2H), 7.33 (d, J = 9 Hz, 2H), 7.77 (d, J = 9 Hz, 2H).

25

Step 10: Preparation of 1-[2-(4-fluorophenyl)-4,4-di(tosylmethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

30

Under nitrogen, a solution of 2.34 mmol of the crude 1-[2-(4-fluorophenyl)-4,4-di(hydroxymethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (prepared in Step 9) in 8 mL of pyridine at ambient temperature was treated with 1.2 g (6.3 mmol) of *p*-toluenesulfonyl chloride (tosyl chloride). The resulting solution was stirred at room temperature for 17 hours, concentrated in vacuo, and chromatographed on silica gel to give 1.06 g (66% overall

35

yield from Step 9) of 1-[2-(4-fluorophenyl)-4,4-di(tosylmethyl)cyclopenten-1-yl]-4-

(methylsulfonyl)benzene as a colorless solid: NMR

(CDCl₃) δ 2.46 (s, 6H), 2.73 (s, 3H), 3.04 (s, 3H), 4.05

5 (s, 4H), 6.85-7.0 (m, 4H), 7.20 (d, J = 8 Hz, 2H), 7.34 (d, J = 8 Hz, 4H), 7.75 (d, J = 8 Hz, 6H).

Step 11: Preparation of 1-[2-(4-fluorophenyl)-4,4-

di(fluoromethyl)cyclopenten-1-yl]-4-

10 (methylsulfonyl)benzene


Under nitrogen, 20 mL (20 mmol) of Bu₄NF (1.0 M in THF) was added to a stirred solution of 3.7 g (5.4 mmol) of 1-[2-(4-fluorophenyl)-4,4-

15 di(tosylmethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (prepared in Step 10) in 10 mL of anhydrous THF. The reaction was stirred at reflux for 14 hours and allowed to cool to ambient temperature. Purification by silica gel chromatography (MPLC) using

20 hexane/ethyl acetate (9:1) gave 1.0 g (49%) of 1-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene as a pale-yellow solid: mp 134-135°C; NMR (CDCl₃) δ 2.84 (s, 4H), 3.04 (s, 3H), 4.52 (d, J = 47 Hz, 2H), 6.95 (t, J = 9 Hz, 2H), 7.06-7.13 (m,

25 2H), 7.32 (d, J = 9 Hz, 2H), 7.79 (d, J = 9 Hz, 2H); HRMS. Calc'd for C₂₀H₁₉F₃O₂S: 380.1058. Found: 380.1077. Anal. Calc'd for C₂₀H₁₉F₃O₂S + 0.30 H₂O: C, 62.26; H, 5.12; F, S, 8.31. Found: C, 62.47; H, 5.26; S, 8.47.

Example 17




5 1-[2-(4-Fluorophenyl)-4,4-
 di(trifluoromethyl)cyclopenten-1-yl]-4-
 (methylsulfonyl)benzene

Following the general procedures outlined in
 10 Synthetic Schemes XV, XX, XXI, and XXII, 1-[2-(4-
 fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]-
 4-(methylsulfonyl)benzene was isolated as a colorless
 solid: mp 137.7-138.5°C; NMR (CDCl₃) δ 3.04 (s, 3H),
 3.36 (s, 4H), 6.98 (t, J= 9 Hz, 2H), 7.07-7.15 (m, 2H),
 15 7.33 (t, J= 9 Hz, 2H), 7.82 (t, J= 9 Hz, 2H); MS (FAB)
 m/e (rel intensity) 459 (100), 374 (10); HRMS. Calc'd
 for C₂₀H₁₅F₇O₂S: 452.0681. Found: 452.0712. Anal.
 Calc'd for C₂₀H₁₅F₇O₂S + 0.23 H₂O: C, 53.16; 3.60; F,
 28.11. Found: C, 52.85; H, 3.40; N, 28.06.

20

Example 18




25 1-[2-(4-Fluorophenyl)-4,4-diethylcyclopenten-1-
 yl]-4-(methylsulfonyl)benzene

Following the general procedures outlined in Synthetic Schemes XV, XX, XXI, and XXII, 1-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-4-

5 (methylsulfonyl)benzene was isolated as a pale-yellow oil: NMR (CDCl₃) δ 0.91 (t, J = 8 Hz, 6H), 1.56 (q, J = 8 Hz, 4H), 2.69 (s, 4H), 3.03 (s, 3H), 6.92 (t, J = 8 Hz, 2H), 7.09 (t, J = 8 Hz, 2H), 7.31 (t, J = 8 Hz, 2H), 7.75 (t, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 372 (100), 343 (46), 264 (18), 235 (20); HRMS. Calc'd for C₂₂H₂₅FO₂S: 372.1559. Found: 372.1532. Anal. Calc'd for C₂₂H₂₅FO₂S: C, 69.08; H, 6.88; F, 4.97; S, 8.38. Found: C, 69.05; H, 6.96; F, 4.92; S, 8.71.

15 Example 19



20 4-[2-(4-Fluorophenyl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 18) was converted to 4-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 100.8-101.2°C; NMR (CDCl₃) δ 0.91 (t, J = 8 Hz, 6H), 1.49-1.60 (m, 4H), 2.68 (s, 4H), 4.69-4.77 (m, 2H), 6.92 (t, J = 9 Hz, 2H), 7.05-7.13 (m, 2H), 7.26 (t, J = 9 Hz, 2H), 7.74 (t, J = 9 Hz, 2H); MS (EI) m/e (rel intensity) 373 (100), 344 (38), 264 (12), 235 (25), 109 (63); HRMS. Calc'd

117


for $C_{21}H_{24}FNO_2S$: 373.1512. Found: 373.1572. Anal.

Calc'd for $C_{21}H_{24}FNO_2S$: C, 67.53; H, 6.48; N, 3.75.

Found: C, 67.55; H, 6.43; N, 3.75.

5

Example 20




10

4-[2-(4-Fluorophenyl)cyclopenten-1-yl]benzenesulfonamide

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 1) was converted to 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 173-174°C; NMR ($CDCl_3$) δ 2.02-2.15 (m, 2H), 2.92 (t, J = 7 Hz, 4H), 4.73 (br s, 2H), 6.93 (t, J = 10 Hz, 2H), 7.06-7.15 (m, 2H), 7.28 (d, J = 8 Hz, 2H), 7.75 (d, J = 8 Hz, 2H); MS (EI) m/e (rel intensity) 317 (100) 237 (40), 221 (30), 109 (55); HRMS. Calc'd for $C_{17}H_{16}FNO_2S$: 317.0886. Found: 317.0916. Anal. Calc'd for $C_{17}H_{16}FNO_2S$: C, 64.28; H, 5.05; N, 4.42. Found: C, 63.98; H, 5.29; N, 4.21.

25

Example 21


5 **4-[2-(4-Chlorophenyl)cyclopenten-1-yl]benzenesulfonamide**

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 3) was converted to 4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 177-178°C; NMR (CDCl₃) δ 2.09 (p, J= 8 Hz, 2H), 2.90 (t, J= 8 Hz, 4H), 4.78 (s, 2H), 7.06 (d, J= 9 Hz, 2H), 7.20 (d, J= 9 Hz, 2H), 7.27 (d, J= 9 Hz, 2H), 7.76 (d, J= 9 Hz, 2H); MS (EI) m/e (rel intensity) 335 (57), 333 (100), 253 (24), 218 (70), 217 (47), 202 (36), 125 (29), 115 (36), 91 (24); HRMS. Calc'd for C₁₇H₁₆ClNO₂S: 333.0590. Found: 333.0587.

15 **Anal.** Calc'd for C₁₇H₁₆ClNO₂S: C, 60.04; H, 4.95; Cl, 10.42; N, 4.12; S, 9.43. Found: C, 59.83; H, 4.84; Cl, 10.99; S, 10.77.

Example 22

25




4-[2-(4-(Trifluoromethyl)phenyl)

cyclopenten-1-yl]benzenesulfonamide

- Following the general procedures outlined in Synthetic Scheme XXIII, 1-[2-(4-
- 5 (trifluoromethyl)phenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 6) was converted to 4-[2-(4-(trifluoromethyl)phenyl)cyclopenten-1-
- 10 yl]benzenesulfonamide as a colorless solid: mp 164-165°C (dec); NMR C(CDCl₃) δ 2.05-2.18 (m, 2H), 2.93 (t, J = 8 Hz, 4H), 4.75 (s, 2H), 7.20-7.30 (m, 4H), 7.48 (t, J = 8 Hz, 2H), 7.66 (t, J = 8 Hz, 2H); MS (FAB) m/e (rel intensity) 392 (26), 374 (100), 334 (13), 308 (8); HRMS. Calc'd for C₁₈H₁₆F₃NO₂S: 367.0854. Found: 367.0832.
- 15 Anal. Calc'd for C₁₈H₁₆F₃NO₂S: C, 58.85; H, 4.39; N, 3.81. Found: C, 58.75; H, 4.45; N, 3.53.


120

Example 23

5 1-[2-(4-Methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

Following the general procedures outlined in Synthetic Schemes I, VII, and X, 1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene was isolated as a colorless solid: mp 128.8-129.8°C; NMR (CDCl₃) δ 2.01-2.14 (m, 2H), 2.90 (t, J= 8 Hz, 4H), 3.04 (s, 3H), 3.79 (s, 3H), 6.78 (t, J= 9 Hz, 2H), 7.08 (t, J= 9 Hz, 2H), 7.36 (t, J= 9 Hz, 2H), 7.75 (t, J= 9 Hz, 2H); MS (FAB) m/e (rel intensity) 335 (100); HRMS. Calc'd for C₁₉H₂₀O₃S: 328.1133. Found: 328.1125. Anal. Calc'd for C₁₉H₂₀O₃S: C, 69.48; H, 6.14; S, 9.76. Found: C, 69.62; H, 6.19; S, 9.80.

20

Example 24

25 4-[2-(4-Methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide


Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(4-methoxyphenyl)

121

cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 23) was converted to 4-[2-(4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 171-175°C (dec); NMR (CDCl₃) δ

- 5 2.00-2.12 (m, 2H), 2.89 (t, J = 8 Hz, 4H), 3.79 (s, 3H),
4.70-4.76 (m, 2H), 6.77 (t, J = 9 Hz, 2H), 7.08 (t, J = 9
Hz, 2H), 7.30 (t, J = 9 Hz, 2H), 7.73 (t, J = 9 Hz, 2H);
MS (FAB) m/e (rel intensity) 329 (100), 297 (11), 249
(64); HRMS. Calc'd for C₁₈H₁₉NO₃S: 329.1086. Found:
10 329.1112. Anal. Calc'd for C₁₈H₁₉NO₃S: C, 65.63; H,
5.81; N, 4.25. Found: C, 65.46; H, 5.89; N, 4.26.

Example 25




15

1-[2-(2,3-Difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

- 20 Following the general procedures outlined in
Synthetic Schemes I, VII, and X, 1-[2-(2,3-
difluorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene was isolated as a colorless
solid: mp 151-152°C; NMR (CDCl₃) δ 2.14 (p, J = 8 Hz,
25 2H), 2.87-3.00 (m, 4H), 3.02 (s, 3H), 6.77-6.84 (m, 1H),
6.91-7.11 (m, 2H), 7.28 (d, J = 9 Hz, 2H), 7.74 (d, J = 9
Hz, 2H); MS (EI) m/e (rel intensity) 334 (100), 255
(54), 227 (24), 128 (29), 127 (33); HRMS. Calc'd for
C₁₈H₁₆F₂O₂S: 334.0839. Found: 334.0835. Anal. Calc'd
30 for C₁₈H₁₆F₂O₂S: C, 64.66; H, 4.82; S, 9.59. Found: C,
64.67; H, 4.87; S, 9.46.

Example 26




5

4-[2-(2,3-Difluorophenyl)cyclopenten-1-yl]benzenesulfonamide

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 25) was converted to 4-[2-(2,3-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 133-134°C; NMR (CDCl₃) δ 2.13 (p, J = 8 Hz, 2H), 2.85-3.00 (m, 4H), 4.75 (s, 2H), 6.76-6.84 (m, 1H), 6.90-7.11 (m, 2H), 7.20-7.27 (m, 2H), 7.72 (d, J = 8 Hz, 2H); HRMS. Calc'd for C₁₇H₁₅F₂NO₂S: 335.0792. Found: 335.0774. Anal. Calc'd for C₁₇H₁₅F₂NO₂S + 0.18 H₂O: C, 60.29; H, 4.57; F, 11.22; N, 4.14; S, 9.47. Found: C, 60.27; H, 4.69; F, 11.35; N, 4.06; S, 9.52.

20

Example 27




25

1-[2-(3,4-Dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

Following the general procedures outlined in Synthetic Schemes I, VII, and X, 1-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene was isolated as a colorless solid: mp 100-101°C; NMR (CDCl₃) δ 2.10 (p, J= 8 Hz, 2H), 2.85-2.96 (m, 4H), 3.05 (s, 3H), 6.91 (dd, J= 9 and 2 Hz, 1H), 7.23-7.28 (m, 2H), 7.32 (d, J= 9 Hz, 2H), 7.79 (d, J= 9 Hz, 2H); HRMS. Calc'd for C₁₈H₁₆Cl₂O₂S: 366.0248. Found: 366.0269. Anal. Calc'd for C₁₈H₁₆Cl₂O₂S: C, 58.86; H, 4.39; Cl, 19.31; S, 8.73. Found: C, 58.53; H, 4.53; Cl, 19.26; S, 8.78.


Example 28



4-[2-(3,4-Dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide


Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 27) was converted to 4-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 134-135°C; NMR (CDCl₃) δ 2.10 (p, J= 8 Hz, 2H), 2.84-2.95 (m, 4H), 4.76 (s, 2H), 6.91 (dd, J= 9 and 2 Hz, 1H), 7.23-7.30 (m, 4H), 7.78 (d, J= 9 Hz, 2H); MS (FAB) m/e 372, 370, 368. Anal. Calc'd for C₁₇H₁₅Cl₂NO₂S: C, 55.43; H, 4.11; Cl, 19.25; N, 3.80; S, 8.71. Found: C, 55.57; H, 4.02; Cl, 19.24; N, 3.67; S, 8.62.

Example 29

5 1-[2-(3-Fluoro-4-methoxyphenyl)cyclopenten-1-yl]-
4-(methylsulfonyl)benzene

Following the general procedures outlined in Synthetic Schemes I, VII, and X, 1-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene was isolated as a colorless solid: mp 127.8-128.6°C; NMR (CDCl₃) δ 2.02-2.14 (m, 2H), 2.84-2.94 (m, 4H), 3.05 (s, 3H), 3.86 (s, 3H), 6.76-6.92 (m 3H), 7.34 (t, J= 9 Hz, 2H), 7.78 (t, J= 9 Hz, 2H); MS (EI) m/e (rel intensity) 346 (100), 315 (7), 267 (10), 191 (10), 139 (12), 107 (7); HRMS. Calc'd for C₁₉H₁₉FO₃S: 346.1039. Found: 346.1049. Anal. Calc'd for C₁₉H₁₉FO₃S: C, 65.88; H, 5.53; F, 5.48. Found: C, 65.71; H, 5.49; F, 5.13.


20 Example 30

25 4-[2-(3-Fluoro-4-methoxyphenyl)cyclopenten-
1-yl]benzenesulfonamide

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]-4-

- (methylsulfonyl)benzene (the title compound of Example 29) was converted to 4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 170.0-170.8°C; NMR (CDCl₃) δ 2.01-2.13 (m, 2H), 2.83-2.93 (m, 4H), 3.87 (s, 3H), 4.75 (s, 2H), 6.76-6.92 (m, 3H), 7.30 (t, *J* = 9 Hz, 2H), 7.76 (t, *J* = 2H); MS (EI) *m/e* (rel intensity) 347 (100), 316 (12), 267 (12), 252 (19), 236 (20), 191 (16), 139 (12); HRMS. Calc'd for C₁₈H₁₈FNO₃S: 347.0991. Found: 347.0955.
- 10 Anal. Calc'd for C₁₈H₁₈FNO₃S: C, 62.23; H, 5.22; N, 4.03. Found: 62.37; H, 5.36; 3.99.


Example 31



15

1-[2-(3-Chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

- 20 Following the general procedures outlined in Synthetic Schemes I, VII, and X, 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene was isolated as a colorless solid: mp 118-120°C; NMR (CDCl₃) δ 2.01-2.14 (m, 2H),
- 25 2.83-2.95 (m, 4H), 3.04 (s, 3H), 3.87 (s, 3H), 6.77 (d, *J* = 9 Hz, 1H), 6.94 (d, *J* = 9 Hz, 1H), 7.18 (d, *J* = 2 Hz, 1H), 7.34 (t, *J* = 9 Hz, 2H), 7.77 (t, *J* = 9 Hz, 2H); MS (EI) *m/e* (rel intensity) 362 (100), 327 (13), 248 (18), 233 (8); HRMS. Calc'd for C₁₉H₁₉ClO₃S: 362.0743.
- 30 Found: 362.0727. Anal. Calc'd for C₁₉H₁₉ClO₃S: C, 62.89; H, 5.28; Cl, 9.77. Found: C, 62.89; H, 5.42; 9.95.


Example 32

5 **4-[2-(3-Chloro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide**

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 31) was converted to 4-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 178.8-180.0°C; NMR (CDCl₃) δ 2.01-2.13 (m, 2H), 2.83-2.93 (m, 4H), 3.87 (s, 3H), 4.75 (s, 2H), 6.76 (d, J= 9 Hz, 1H), 6.94 (dd, J= 9 and 2 Hz, 1H), 7.20 (d, J= 2 Hz, 1H), 7.30 (t, J= 8 Hz, 2H), 7.76 (t, J= 8 Hz, 2H); MS (EI) m/e (rel intensity) 363 (100), 328 (14), 268 (16), 233 (17); HRMS. Calc'd for C₁₈H₁₈ClNO₃S: 363.0696. Found: 363.0701. Anal. Calc'd for C₁₈H₁₈ClNO₃S: C, 59.41; H, 5.00; N, 3.85. Found: C, 59.12; H, 5.06; N, 3.69.

Example 33

25




1-[2-(3-Chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

Following the general procedures outlined in Synthetic Schemes I, VII, and X, 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene was isolated as a colorless solid: mp 87.5-88.5°C; NMR (CDCl₃) δ 2.03-2.15 (m, 2H), 2.84-2.95 (m, 4H), 3.04 (s, 3H), 3.87 (s, 3H), 6.91-7.02 (m, 2H), 7.19 (dd, J= 8 and 2 Hz, 1H), 7.31 (t, J= 9 Hz, 2H), 7.79 (t, J= 9 Hz, 2H); MS (EI) m/e (rel intensity) 350 (100), 315 (6), 271 (22), 236 (69); HRMS. Calc'd for C₁₈H₁₆ClFO₂S: 350.0544. Found: 350.0523. Anal. Calc'd for C₁₈H₁₆ClFO₂S: C, 61.62; H, 4.60; F, 5.42. Found: C, 61.37; H, 4.67; F, 5.09.

15


Example 34



20

4-[2-(3-Chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 33) was converted to 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 150.5-151.5°C; NMR (CDCl₃) δ 2.02-2.16 m, 2H), 2.83-2.96 (m, 4H), 4.78 (s, 2H), 6.90-7.03 (m, 2H), 7.18-7.24 (m, 1H), 7.28 (t, J= 9 Hz, 2H), 7.78 (t, J= 9 Hz, 2H); MS (EI) m/e (rel intensity) 351 (100), 316 (16), 271 (40), 236 (28); HRMS. Calc'd for C₁₇H₁₅ClFNO₂S: 351.0496. Found: 351.0516. Anal. Calc'd for C₁₇H₁₅ClFNO₂S: C, 58.04; H, 4.30; N, 3.98. Found: C, 57.94; H, 4.41; N, 3.93.


Example 35

5

5-[2-[4-(Methylsulfonyl)phenyl]cyclopenten-1-yl]-1,3-benzodioxole

Following the general procedures outlined in Synthetic Schemes I, VII, and X, 5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]-1,3-benzodioxole was isolated as a colorless solid: mp 140.0-140.7°C; NMR (CDCl₃) δ 2.06 (p, J = 8 Hz, 2H), 2.82-2.93 (m, 4H), 3.04 (s, 3H), 5.93 (s, 2H), 6.59-6.72 (m, 3H), 7.35 (d, J = 9 Hz, 2H), 7.76 (d, J = 9 Hz, 2H); HRMS. Calc'd for C₁₉H₁₈O₄S: 342.0926. Found: 342.0932. Anal. Calc'd for C₁₉H₁₈O₄S: C, 66.65; H, 5.30. Found: C, 66.43; H, 5.47.

20

Example 36


25

4-[2-(2-Methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide

Following the general procedure outlined in Synthetic Scheme XI (with the substitution of 33 for

43), 1-(2-bromocyclopenten-1-yl)-4-(methylsulfonyl)benzene (prepared in Step 1 of Example 11) was reacted with 2-methylpyridin-5-yltrimethyltin (prepared in Synthetic Scheme VIII) to provide 4-[2-(2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 108.0-109.5°C; NMR (CDCl₃) δ 2.05-2.19 (m, 2H), 2.46 (s, 3H), 2.86 (t, J = 7 Hz, 4H), 3.05 (s, 3H), 7.02 (d, J = 8 Hz, 1H), 7.32 (br s, 1H), 7.33 (d, J = 8 Hz, 2H), 7.78 (d, J = 8 Hz, 2H), 8.82 (d, J = 2Hz, 1H); MS (CI) m/e (rel intensity) 314 (100), 234 (5), 89 (20); HRMS. Calc'd for C₁₈H₁₉LiNO₂S: 320.1297. Found: 320.1281. Anal. Calc'd for C₁₈H₁₉NO₂S: C, 69.01; H, 6.07; N, 4.47. Found: C, 68.91; H, 6.40; 4.28.

Example 37



1-[2-(4-Fluorophenyl)cyclopenten-1-yl]-4-(fluoromethylsulfonyl)benzene


Under nitrogen, 2.7 mL (3.8 mmol) of methyllithium (1.4 M in ether) was added to a stirred solution of 1.0 g (3.2 mmol) of 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 1) in 10 mL of anhydrous THF at -78°C. The reaction was allowed to warm to ambient temperature and stir for 1 hour. A solution of 1.2 g (4.8 mmol) of N-fluorodibenzenesulfonamide (Aldrich) in 5 mL of anhydrous THF was slowly added at 0°C. The reaction was stirred for 1 hour at ambient temperature and quenched with water. Purification by

130

silica gel chromatography (MPLC) using hexane/ethyl acetate (9:1) gave 110 mg (10%) of 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-

- (fluoromethylsulfonyl)benzene as a colorless solid: mp 123-124°C; NMR (CDCl₃) δ 2.10 (p, J= 7 Hz, 2H), 2.92 (t, J= 7 Hz, 4H), 5.11 (d, J= 47 Hz, 2H), 6.94 (dd, J= 9 and 2 Hz, 2H), 7.07-7.14 (m, 2H), 7.36 (d, J= 9 Hz, 2H), 7.77 (d, J= 9 Hz, 2H); MS (EI) m/e (rel intensity) 334 (100), 237 (24), 209 (22); HRMS. Calc'd for C₁₈H₁₆F₂O₂S: 334.0839. Found: 334.0828. Anal. Calc'd for C₁₈H₁₆F₂O₂S + 0.22 H₂O: C, 63.90; H, 4.90; F, 11.23; S, 9.48. Found: C, 63.89; H, 4.90; F, 11.07; S, 9.76.

Example 38




15

1-[2-(3,4-Difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene

- Following the general procedures outlined in Synthetic Schemes I, VII, and X, 1-[2-(3,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene was isolated as a colorless solid: mp 114-115 °C; NMR (CDCl₃) δ 2.10 (p, J= 8 Hz, 2H), 2.85-2.96 (m, 4H), 3.05 (s, 3H), 6.79-6.86 (m, 1H), 6.89-7.07 (m, 2H), 7.32 (d, J= 9 Hz, 2H), 7.79 (d, J= 9 Hz, 2H); MS (FAB) m/e 341 (M+Li); HRMS. Calc'd for C₁₈H₁₇F₂O₂S: 335.0917. Found: 335.0925. Anal. Calc'd for C₁₈H₁₆F₂O₂S: C, 64.66; H, 4.82; F, 11.36. Found: C, 64.77; H, 4.91; F, 11.39.

Example 39



4-[2-(3,4-Difluorophenyl)cyclopenten-1-yl]benzenesulfonamide

5

Following the general procedure outlined in Synthetic Scheme XXIII, 1-[2-(3,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene (the title compound of Example 38) was converted to 4-[2-(3,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide as a colorless solid: mp 147-147.5 °C; NMR (CDCl₃) δ 2.10 (p, J = 9 Hz, 2H), 2.84-2.95 (m, 4H), 4.76 (s, 2H), 6.80-6.86 (m, 1H), 6.90-7.07 (m, 2H), 7.27 (d, J = 9 Hz, 2H), 7.78 (d, J = 9 Hz, 2H); MS (FAB) m/e 342 (M+Li); HRMS. Calc'd for C₁₇H₁₆F₂NO₂S: 336.0870. Found: 336.0856. Anal. Calc'd for C₁₇H₁₅F₂NO₂S: C, 60.88; H, 4.51; N, 4.18. Found: C, 60.97; H, 4.56; N, 4.15.

10

15

BIOLOGICAL EVALUATION

20 Rat Carrageenan Foot Pad Edema Test

The carrageenan foot edema test was performed with materials, reagents and procedures essentially as described by Winter, et al., (Proc. Soc. Exp. Biol. Med., 111, 544 (1962)). Male Sprague-Dawley rats were selected in each group so that the average body weight was as close as possible. Rats were fasted with free access to water for over sixteen hours prior to the test. The rats were dosed orally (1 mL) with compounds suspended in vehicle containing 0.5% methylcellulose and .025% surfactant, or with vehicle alone. One hour later

25

30

a subplantar injection of 0.1 mL of 1% solution of carrageenan/sterile 0.9% saline was administered and the volume of the injected foot was measured with a displacement plethysmometer connected to a pressure transducer with a digital indicator. Three hours after the injection of the carrageenan, the volume of the foot was again measured. The average foot swelling in a group of drug-treated animals was compared with that of a group of placebo-treated animals and the percentage inhibition of edema was determined (Otterness and Bliven, Laboratory Models for Testing NSAIDs, in Non-steroidal Anti-Inflammatory Drugs, (J. Lombardino, ed. 1985)). Results are shown in Table I.

15

Rat Carrageenan-induced Analgesia Test

The analgesia test using rat carrageenan was performed with materials, reagents and procedures essentially as described by Hargreaves, et al., (Pain, 32, 77 (1988)). Male Sprague-Dawley rats were treated as previously described for the Carrageenan Foot Pad Edema test. Three hours after the injection of the carrageenan, the rats were placed in a special plexiglass container with a transparent floor having a high intensity lamp as a radiant heat source, positionable under the floor. After an initial twenty minute period, thermal stimulation was begun on either the injected foot or on the contralateral uninjected foot. A photoelectric cell turned off the lamp and timer when light was interrupted by paw withdrawal. The time until the rat withdraws its foot was then measured. The withdrawal latency in seconds was determined for the control and drug-treated groups, and percent inhibition of the hyperalgesic foot withdrawal determined. Results are shown in Table I.

TABLE I.

	RAT PAW EDEMA		ANALGESIA	
	% Inhibition		% Inhibition	
5	@ 10mg/kg body weight		@ 20mg/kg body weight	
	Examples			
	1	32	66	
	2	39		
	3	33	41	
10	4	25		
	5	16		
	6	39	29	
	11	28	39	
	12	45 ¹		
15	13	34 ²	14 ²	
	14	31 ²		
	15	41 ²		
	20	26	44 ²	
	21	49 ²		
20	23	22	15 ²	
	25	32 ²		
	28	10	26 ²	
	30	47 ²		
	31	28 ²		
25	32	13	17 ²	
	34	37	46 ²	
	35	6 ²		
	36	33 ²		
30	1 - Assay performed at 20 mg/kg body weight.			
	2 - Assay performed at 30 mg/kg body weight.			

Evaluation of COX I and COX II activity in vitro

35 The compounds of this invention exhibited inhibition in vitro of COX II. The COX II inhibition activity of the compounds of this invention illustrated in the Examples was determined by the following methods.

a. Preparation of recombinant COX baculoviruses

A 2.0 kb fragment containing the coding region
5 of either human or murine COX-I or human or murine COX-II was cloned into a BamHI site of the baculovirus transfer vector pVL1393 (Invitrogen) to generate the baculovirus transfer vectors for COX-I and COX-II in a manner similar to the method of D.R. O'Reilly et al
10 (*Baculovirus Expression Vectors: A Laboratory Manual* (1992)). Recombinant baculoviruses were isolated by transfecting 4 µg of baculovirus transfer vector DNA into SF9 insect cells (2×10^8) along with 200 ng of linearized baculovirus plasmid DNA by the calcium
15 phosphate method. See M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agric. Exp. Station Bull. 1555 (1987). Recombinant viruses were purified by three rounds of plaque purification and high titer (10^7 -
20 10^8 pfu/ml) stocks of virus were prepared. For large scale production, SF9 insect cells were infected in 10 liter fermentors (0.5×10^6 /ml) with the recombinant baculovirus stock such that the multiplicity of infection was 0.1. After 72 hours the cells were
25 centrifuged and the cell pellet homogenized in Tris/Sucrose (50 mM: 25%, pH 8.0) containing 1% 3-[(3-cholamidopropyl)dimethylammonio] -1-propanesulfonate (CHAPS). The homogenate was centrifuged at 10,000xG for 30 minutes, and the resultant supernatant was stored at
30 -80°C before being assayed for COX activity.

b. Assay for COX I and COX II activity:

COX activity was assayed as PGE₂ formed/µg
35 protein/time using an ELISA to detect the prostaglandin released. CHAPS-solubilized insect cell membranes containing the appropriate COX enzyme were incubated in a potassium phosphate buffer (50 mM, pH 8.0) containing

epinephrine, phenol, and heme with the addition of arachidonic acid (10 μM). Compounds were pre-incubated with the enzyme for 10-20 minutes prior to the addition of arachidonic acid. Any reaction between the

5 arachidonic acid and the enzyme was stopped after ten minutes at 37°C/room temperature by transferring 40 μl of reaction mix into 160 μl ELISA buffer and 25 μM indomethacin. The PGE₂ formed was measured by standard ELISA technology (Cayman Chemical). Results are shown in

10 Table II.

TABLE II.

		HUMAN COX I ID50 μM	HUMAN COX II ID50 μM
	Examples		
	1	>100	<.1
	2	>100	<.1
20	3	>100	<.1
	4	>100	<.1
	5	>100	<.1
	6	>100	.9
	7	>100	2.2
25	8	>100	77.9
	9	>100	3.2
	10	>100	6.6
	11	>100	.2
	12	18.3	<.1
30	13	.5	<.1
	14	1.6	<.1
	15	<.1	<.1
	16	58	<.1
	17	>100	<.1
35	18	>100	65
	19	10	3.9
	20	4.2	<.1
	21	1.3	<.1

TABLE II. (cont.)

		HUMAN COX I ID50 μ M	HUMAN COX II ID50 μ M
5	Examples		
	22	4.0	.1
	23	9.9	<.1
	24	.2	<.1
10	25	>100	<.1
	26	3.6	<.1
	27	>100	<.1
	28	3.7	<.1
	29	>100	.1
15	30	8.1	<.1
	31	>100	.1
	32	21.3	<.1
	33	>100	<.1
	34	2.1	<.1
20	35	>100	<.1
	36	>100	.8
	37	>100	<.1
	38	>100	1.0
	39	9.9	<.1
25			

Also embraced within this invention is a class of pharmaceutical compositions comprising one or more compounds of Formula I in association with one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The compounds and composition may, for example, be administered intravascularly,

intraperitoneally, subcutaneously, intramuscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier.

The amount of therapeutically active compound that is administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the disease, the route and frequency of administration, and the particular compound employed, and thus may vary widely. The pharmaceutical compositions may contain active ingredient in the range of about 0.1 to 1000 mg, preferably in the range of about 0.5 to 500 mg and most preferably between about 1 and 100 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between about 0.1 and about 50 mg/kg body weight and most preferably from about 1 to 20 mg/kg body weight, may be appropriate. The daily dose can be administered in one to four doses per day.


For therapeutic purposes, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl

esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.


Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

What is claimed is:

1. A compound of Formula I




wherein A is selected from



wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

wherein each of R³ through R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl;

provided that when A is



R⁵ cannot be hydrido when R¹⁰ is selected from hydrido, cyano, halo, methyl, trifluoromethyl and methoxy; further provided that R¹⁰ cannot be hydrido when R⁵ is selected from cyano, halo, methyl, trifluoromethyl and methoxy; and further provided that R⁵ and R¹⁰ are not both methoxy; or a pharmaceutically suitable salt thereof.


2. Compound of Claim 1 wherein each of R^1 and R^2 is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxy carbonyl and carboxyl; and wherein each of R^3 through R^{12} is
5 independently selected from hydrido, halo, lower alkyl, lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof.

10

3. Compound of Claim 2 wherein each of R^1 and R^2 is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo,
15 fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R^3 through R^{12} is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl,
25 isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl,
35 methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof.

141

4. A compound of Formula II



II

5 wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

 wherein each of R³ through R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, 10 haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl;

 provided that R⁵ cannot be hydrido when R¹⁰ is selected from hydrido, cyano, halo, methyl, trifluoromethyl and methoxy; further provided that R¹⁰ cannot be hydrido 15 when R⁵ is selected from cyano, halo, methyl, trifluoromethyl and methoxy; and further provided that R⁵ and R¹⁰ are not both methoxy; or a pharmaceutically suitable salt thereof.

20 5. Compound of Claim 4 wherein each of R¹ and R² is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxycarbonyl and carboxyl; and wherein each of R³ through R¹² is independently selected from hydrido, halo, lower alkyl, 25 lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof.

30

 6. The compound of Claim 5 wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl,

propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, 5 pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, 10 pentoxycarbonyl and carboxyl; and wherein each of R³ through R¹² is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, 15 chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, 20 trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl, methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof.

25

7. Compound of Claim 6 selected from compounds, and their pharmaceutically-acceptable salts, of the group consisting of

- 30 1-(2-phenylcyclopenten-1-yl)-4-(methylsulfonyl)benzene,
1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
35 1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-

- (methylsulfonyl)benzene;
1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-cyanophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
5 1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
10 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(2,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
15 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
20 4-[2-(2,3-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
30 4-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
35 1-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide;

- 1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
5 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(fluoromethylsulfonyl)benzene;
1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(fluoromethylsulfonyl)benzene;
10 5-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
1-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
15 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
20 1-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
30 4-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
1-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
35 4-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]
benzenesulfonamide;
- 1-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 5 1-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-
10 (methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3,4-difluorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 15 1-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-
20 (methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 25 1-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-
30 (methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 35 1-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;

- 1-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-
4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 5 1-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-
10 (methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]-
4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 15 1-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-
20 (methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-[3,4-di(diifluoromethyl)phenyl]cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 25 1-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]-4-
30 (methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 4-[2-(4-methylthiophenyl)cyclopenten-1-
yl]benzenesulfonamide;
- 35 4-[2-(4-cyanophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-hydroxymethylphenyl)cyclopenten-1-
yl]benzenesulfonamide;

- 4-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 5 4-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 10 5-[2-(4-(aminosulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
- 4-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-phenylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 30 4-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 5 4-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 30 4-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-[3,4-di(diifluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;

4-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;

4-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide; and

5 4-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide.

8. Compound of Claim 6 which is 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene,
10 or a pharmaceutically-acceptable salt thereof.

9. Compound of Claim 6 which is 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene, or a pharmaceutically-acceptable
15 salt thereof.

10. Compound of Claim 6 which is 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide, or a
pharmaceutically-acceptable salt thereof.

20

11. Compound of Claim 6 which is 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable
salt thereof.


25

12. Compound of Claim 6 which is 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene,
or a pharmaceutically-acceptable salt thereof.

30

13. Compound of Claim 6 which is 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide,
or a pharmaceutically-acceptable salt thereof.

14. A compound of Formula III



5

wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl;

wherein R⁵ is selected from alkylsulfonyl, haloalkylsulfonyl and sulfamyl; and


wherein B is phenyl or pyridyl, wherein B is optionally substituted at a substitutable position with alkyl, halo, alkylthio, cyano, haloalkyl, alkoxy, hydroxyalkyl and alkoxyalkyl; or a pharmaceutically-acceptable salt thereof.

15. The compound of Claim 14 wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; wherein R⁵ is selected from methylsulfonyl, trifluoromethylsulfonyl and sulfamyl; and

wherein B is optionally substituted with a radical select d from fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl,

hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl and pentoxymethyl; or a pharmaceutically-acceptable salt thereof.

16. A compound of Formula IV



IV

15

wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and
 wherein each of R³ through R⁷ and R⁹ through R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl;
 or a pharmaceutically-acceptable salt thereof.

17. The compound of Claim 16 wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl,

pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, 5 propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R³ through R⁷ and R⁹ through R¹² is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, 10 pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, 15 dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl, 20 methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof.


18. Compound of Claim 17 selected from compounds, and their pharmaceutically-acceptable salts, of 25 the group consisting of

5-fluoro-2-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
5-chloro-2-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
30 5-methyl-2-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
35 4-[2-(5-fluoropyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-chloropyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(5-methylpyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-trifluoromethylpyridin-2-yl)cyclopenten-1-yl]benzenesulfonamide;
5 5-fluoro-2-[2-[4(methylsulfonyl)phenyl]-4-carboxycyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4(methylsulfonyl)phenyl]-4-carboxycyclopenten-1-yl]pyridine;
4-[2-(5-fluoropyridin-2-yl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(5-trifluoromethylpyridin-2-yl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
5-fluoro-2-[2-[4-methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
15 5-fluoro-2-[2-[4-methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
5-trifluoromethyl-2-[2-[4-methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
20 4-[2-(5-fluoropyridin-2-yl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-fluoropyridin-2-yl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(5-trifluoromethylpyridin-2-yl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
2-[2-(4-fluorophenyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
30 2-[2-(4-chlorophenyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-methylphenyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
35 2-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
2-[2-(4-fluorophenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;

- 2-[2-(4-chlorophenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
- 2-[2-(4-methylphenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
- 5 2-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]pyridine-5-sulfonamide;
- 2-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
- 2-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
- 10 2-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]pyridine-5-sulfonamide;
- 2-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]pyridine-5-sulfonamide;
- 15 2-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
- 2-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
- 2-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
- 20 2-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]-5-(methylsulfonyl)pyridine;
- 2-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]pyridine-5-sulfonamide;
- 25 2-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]pyridine-5-sulfonamide;
- 2-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]pyridine-5-sulfonamide; and
- 2-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]pyridine-5-sulfonamide.
- 30

19. A compound of Formula V



V

5 wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

 wherein each of R³ through R⁷, R⁸, R¹⁰, R¹¹ and R¹² is independently selected from hydrido, halo, alkyl, 10 alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl;

 or a pharmaceutically-acceptable salt thereof.

15 20. The compound of Claim 19 wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, 20 chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, 25 propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R³ through R⁸ and R¹⁰ through R¹² is independently selected from hydrido, fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, 30 pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl,

- heptafluoropropyl, difluorochloromethyl,
dichlorofluoromethyl, difluoroethyl, difluoropropyl,
dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy,
butoxy, isopropoxy, tert-butoxy, methylenedioxy,
5 trifluoromethoxy, hydroxymethyl, methoxymethyl,
ethoxymethyl, isopropoxymethyl, tert-butoxymethyl,
propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl,
methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and
sulfamyl; or a pharmaceutically suitable salt thereof.
- 10
21. Compound of Claim 20 selected from
compounds, and their pharmaceutically-acceptable salts, of
the group consisting of
- 15 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-
yl]pyridine;
2-chloro-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-
yl]pyridine;
2-methyl-5-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-
20 yl]pyridine;
2-trifluoromethyl-5-[2-[4-
(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
4-[2-(2-fluoropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
25 4-[2-(2-chloropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
4-[2-(2-methylpyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
4-[2-(2-trifluoromethylpyridin-5-yl)cyclopenten-1-
30 yl]benzenesulfonamide;
2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4-
carboxycyclopenten-1-yl]pyridine;
2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4-
carboxycyclopenten-1-yl]pyridine;
35 4-[2-(2-fluoropyridin-5-yl)-4-carboxycyclopenten-1-yl]
benzenesulfonamide;
4-[2-(2-trifluoromethylpyridin-5-yl)-4-carboxycyclopenten-
1-yl]benzenesulfonamide;

- 2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
2-fluoro-5-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
5 2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
2-trifluoromethyl-5-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
4-[2-(2-fluoropyridin-5-yl)-4,4-dimethylcyclopenten-1-yl]
10 benzenesulfonamide;
4-[2-(2-fluoropyridin-5-yl)-4,4-difluorocyclopenten-1-yl]
benzenesulfonamide;
4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(2-trifluoromethylpyridin-5-yl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
5-[2-(4-fluorophenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
5-[2-(4-chlorophenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
20 5-[2-(4-methylphenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
5-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
25 5-[2-(4-fluorophenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
5-[2-(4-chlorophenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
5-[2-(4-methylphenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
30 5-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]pyridine-2-sulfonamide;
5-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
35 5-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
5-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]pyridine-2-sulfonamide;

- 5-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5 5-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 5-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]-2-(methylsulfonyl)pyridine;
- 10 5-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]pyridine-2-sulfonamide;
- 15 5-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]pyridine-2-sulfonamide;
- 5-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]pyridine-2-sulfonamide.
- 1-[2-(2,3-dimethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(3-fluoro-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(3-trifluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2,3-difluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(3-methyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-trifluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(2,3-dichloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(3-fluoro-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(2,3-dimethoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(3-fluoro-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-[2,3-di(trifluoromethyl)pyridin-5-yl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(3-fluoro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(3-difluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-difluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-2-methoxypyridin-5-yl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;


- 1-[2-[2,3-di(difluoromethyl)pyridin-5-yl]cyclopenten-1-yl]-
4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-2-difluoromethylpyridin-5-yl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(3-fluoro-2-difluoromethylpyridin-5-yl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-2-difluoromethylpyridin-5-yl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(3-chloro-2-difluoromethylpyridin-5-yl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
- 4-[2-(2,3-dimethylpyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-fluoro-2-methylpyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 15 4-[2-(3-chloro-2-methylpyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-methylpyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 20 4-[2-(2,3-difluoropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-methyl-2-fluoropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 25 4-[2-(3-chloro-2-fluoropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-fluoropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 30 4-[2-(2,3-dichloropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-methyl-2-chloropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 35 4-[2-(3-fluoro-2-chloropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-
yl]benzenesulfonamide;

- 4-[2-(3-methoxy-2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2,3-dimethoxy-pyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 5 4-[2-(3-methyl-2-methoxy-pyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-fluoro-2-methoxy-pyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-2-methoxy-pyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(3-chloro-2-methoxy-pyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-[2,3-di(trifluoromethyl)pyridin-5-yl]cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(3-methyl-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-fluoro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(3-chloro-2-trifluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(3-difluoromethyl-2-fluoropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-2-chloropyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-difluoromethyl-2-methoxy-pyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 30 4-[2-[2,3-di(difluoromethyl)pyridin-5-yl]cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-(3-fluoro-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide; and

4-[2-(3-chloro-2-difluoromethylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide.

22. A compound of Formula VI

5



VI

wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxy carbonyl and carboxyl;

wherein R⁵ is selected from alkylsulfonyl, haloalkylsulfonyl and sulfamyl; and

wherein each of R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹¹ and R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl and alkoxyalkyl;

or a pharmaceutically-acceptable salt thereof.

23. The compound of Claim 22 wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; wherein R⁵ is selected from methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl; and wherein each of R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹¹ and R¹² is independently selected from hydrido, fluoro,

chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, 5 pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy, isopropoxy, tert-butoxy, methylenedioxy, trifluoromethoxy, hydroxymethyl, methoxymethyl, 10 ethoxymethyl, isopropoxymethyl, tert-butoxymethyl, propoxyethyl, butoxymethyl, isobutoxymethyl, pentoxymethyl, methylsulfonyl, ethylsulfonyl, trifluoromethylsulfonyl and sulfamyl;


or a pharmaceutically-acceptable salt thereof.

15


24. Compound of Claim 23 selected from compounds, and their pharmaceutically-acceptable salts, of the group consisting of

- 20 4-[2-[4-(methylsulfonyl)phenyl]cyclopenten-1-yl]pyridine;
4-[2-(4-pyridinyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-[4-(methylsulfonyl)phenyl]-4,4-dimethylcyclopenten-1-yl]pyridine;
4-[2-[4-(methylsulfonyl)phenyl]-4,4-di(hydroxymethyl)
25 cyclopenten-1-yl]pyridine;
4-[2-[4-(methylsulfonyl)phenyl]-4,4-difluorocyclopenten-1-yl]pyridine;
4-[2-[4-(methylsulfonyl)phenyl]-4,4-di(fluoromethyl)cyclopenten-1-yl]pyridine;
30 4-[2-[4-(pyridinyl)phenyl]-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-[4-(pyridinyl)phenyl]-4,4-di(hydroxymethyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-[4-(pyridinyl)phenyl]-4,4-difluorocyclopenten-1-yl]benzenesulfonamide; and
35 4-[2-[4-(pyridinyl)phenyl]-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide.

25. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said compound selected from a compound of Formula I




wherein A is selected from



wherein each of R¹ and R² is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

wherein each of R³ through R¹² is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl; or a pharmaceutically-acceptable salt thereof.

26. A pharmaceutical composition comprising a therapeutically-effective amount of Formula II



wherein each of R^1 and R^2 is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

5 wherein each of R^3 through R^{12} is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof.

10 27. A pharmaceutical composition of Claim 26 wherein each of R^1 and R^2 is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxycarbonyl and carboxyl; and wherein each of R^3 through R^{12} is independently selected from
15 hydrido, halo, lower alkyl, lower alkylthio, cyano, lower haloalkyl, lower alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof.

20

28. A pharmaceutical composition of Claim 27 wherein each of R^1 and R^2 is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl,
25 fluoro, chloro, bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl,
30 dichloroethyl, dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R^3 through R^{12} is independently selected from hydrido,
35 fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl,

trifluoromethyl, chloromethyl, dichloromethyl,
trichloromethyl, pentafluoroethyl, heptafluoropropyl,
difluorochloromethyl, dichlorofluoromethyl,
difluoroethyl, difluoropropyl, dichloroethyl,
5 dichloropropyl, methoxy, ethoxy, propoxy, butoxy,
isopropoxy, tert-butoxy, methylenedioxy,
trifluoromethoxy, hydroxymethyl, methoxymethyl,
ethoxymethyl, isopropoxymethyl, tert-butoxymethyl,
propoxyethyl, butoxymethyl, isobutoxymethyl,
10 pentoxymethyl, methylsulfonyl, ethylsulfonyl,
trifluoromethylsulfonyl and sulfamyl; or a
pharmaceutically suitable salt thereof.

29. A pharmaceutical composition of Claim 28
15 wherein compounds, and their pharmaceutically-acceptable
salts, of the group consisting of

1-(2-phenylcyclopenten-1-yl)-4-(methylsulfonyl)benzene,
1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-
20 (methylsulfonyl)benzene;
1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
25 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-cyanophenyl)cyclopenten-1-yl]-4-
30 (methylsulfonyl)benzene;
1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
35 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;

- 1-[2-(2,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
5 1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(2,3-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(2,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
20 1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]benzenesulfonamide;
25 1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
30 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(fluoromethylsulfonyl)benzene;
1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(fluoromethylsulfonyl)benzene;
35 5-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
1-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
- 1-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 20 1-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 4-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
- 1-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]-4-
- 10 (methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]-4-
- 20 (methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]-4-
- 30 (methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-[3,4-di(diifluoromethyl)phenyl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 4-[2-(4-methylthiophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(4-cyanophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 30 4-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 5-[2-(4-(aminosulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
- 35 4-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
5 4-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-phenylcyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(3,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
30 4-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
35 4-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;
5 4-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-[3,4-di(diifluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
25 4-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide; and
4-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide.
30

30. The pharmaceutical composition of Claim 28 wherein 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene; or a pharmaceutically-acceptable salt thereof.

35

31. The pharmaceutical composition of Claim 28 wherein said compound is 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene, or a pharmaceutically-acceptable salt thereof.


32. The pharmaceutical composition of Claim 28 wherein said compound is 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide; or a pharmaceutically-acceptable salt thereof.

33. The pharmaceutical composition of Claim 28 wherein said compound is 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide; or a pharmaceutically-acceptable salt thereof.

34. The pharmaceutical composition of Claim 28 wherein said compound is 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene, or a pharmaceutically-acceptable salt thereof.


35. The pharmaceutical composition of Claim 28 wherein said compound is 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.

36. A method of treating inflammation or an inflammation-associated disorder in a subject, said method comprising administering to the subject having such inflammation or inflammation-associated disorder, a therapeutically-effective amount of a compound of Formula I



I

wherein A is selected from




5

wherein each of R^1 and R^2 is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

wherein each of R^3 through R^{12} is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, haloalkylsulfonyl and sulfamyl; or a pharmaceutically-acceptable salt thereof.

37. A method of treating inflammation or an inflammation-associated disorder in a subject, said method comprising administering to the subject having such inflammation or inflammation-associated disorder, a therapeutically-effective amount of a compound of Formula II



II

wherein each of R^1 and R^2 is independently selected from alkyl, hydrido, hydroxyalkyl, halo, haloalkyl, alkoxycarbonyl and carboxyl; and

wherein each of R^3 through R^{12} is independently selected from hydrido, halo, alkyl, alkylthio, cyano, haloalkyl, alkoxy, haloalkoxy, hydroxyalkyl, alkoxyalkyl,

alkylsulfonyl, haloalkylsulfonyl and sulfamyl; or a pharmaceutically-acceptable salt thereof.

38. The method of of Claim 37 wherein each of
5 R¹ and R² is independently selected from lower alkyl, hydrido, lower hydroxyalkyl, halo, lower haloalkyl, lower alkoxy carbonyl and carboxyl; and wherein each of R³ through R¹² is independently selected from hydrido, halo, lower alkyl, lower alkylthio, cyano, lower haloalkyl,
10 lower alkoxy, lower haloalkoxy, lower hydroxyalkyl, lower alkoxyalkyl, lower alkylsulfonyl, lower haloalkylsulfonyl and sulfamyl; or a pharmaceutically suitable salt thereof;
or a pharmaceutically-acceptable salt or prodrug thereof.

15

39. The method of Claim 38 wherein each of R¹ and R² is independently selected from hydrido, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, hydroxymethyl, fluoro, chloro,
20 bromo, iodo, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl,
25 dichloropropyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tertbutoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl and carboxyl; and wherein each of R³ through R¹² is independently selected from hydrido, fluoro, chloro,
30 bromo, iodo, methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl, methylthio, cyano, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl,
35 difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, butoxy,

isopropoxy, tert-butoxy, methylenedioxy,
trifluoromethoxy, hydroxymethyl, methoxymethyl,
ethoxymethyl, isopropoxymethyl, tert-butoxymethyl,
propoxyethyl, butoxymethyl, isobutoxymethyl,
5 pentoxymethyl, methylsulfonyl, ethylsulfonyl,
trifluoromethylsulfonyl and sulfamyl; or a
pharmaceutically suitable salt thereof.

40. The method of Claim 39 wherein compounds, and
10 their pharmaceutically-acceptable salts, of the group
consisting of

1-(2-phenylcyclopenten-1-yl)-4-(methylsulfonyl)benzene,
1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-
15 (methylsulfonyl)benzene;
1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
20 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-cyanophenyl)cyclopenten-1-yl]-4-
25 (methylsulfonyl)benzene;
1-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
30 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(2,4-difluorophenyl)cyclopenten-1-yl]-4-
35 (methylsulfonyl)benzene;
1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;

- 1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-
4-(methylsulfonyl)benzene;
4-[2-(2,3-difluorophenyl)cyclopenten-1-yl]
benzenesulfonamide;
- 5 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)cyclopenten-1-
yl]benzenesulfonamide;
- 10 4-[2-(2,4-dimethylphenyl)cyclopenten-1-
yl]benzenesulfonamide;
4-[2-(2,4-difluorophenyl)cyclopenten-1-
yl]benzenesulfonamide;
4-[2-(2,4-dichlorophenyl)cyclopenten-1-
yl]benzenesulfonamide;
- 15 1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
- 20 4-[2-(4-fluorophenyl)-4,4-diethylcyclopenten-1-yl]
benzenesulfonamide;
1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-
(methylsulfonyl)benzene;
4-[2-(4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-
yl]benzenesulfonamide;
1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-
(fluoromethylsulfonyl)benzene;
1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-
(fluoromethylsulfonyl)benzene;
- 30 5-[2-(4-(methylsulfonyl)phenyl)cyclopenten-1-yl]-1,3-
benzodioxole;
1-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-
yl]-4-(methylsulfonyl)benzene;
1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-
(methylsulfonyl)benzene;

- 1-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;
5 1-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-fluorophenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(4-trifluoromethylphenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;
4-[2-(4-trifluoromethylphenyl)-4,4-difluorocyclopenten-1-yl]benzenesulfonamide;
15 1-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]-4-(methylsulfonyl)benzene;
4-[2-(4-fluorophenyl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(4-trifluoromethylphenyl)-4-carboxycyclopenten-1-yl]benzenesulfonamide;
1-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
25 1-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
30 1-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3,4-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
35 1-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]-4-
- 10 (methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 1-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]-4-
- 20 (methylsulfonyl)benzene;
- 1-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 25 1-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-
- 30 (methylsulfonyl)benzene;
- 1-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;

- 1-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 5 1-[2-[3,4-di(diifluoromethyl)phenyl]cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 10 1-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 15 4-[2-(4-methylthiophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-cyanophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-hydroxymethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(4-methoxymethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(2,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(4-fluorophenyl)-4,4-di(fluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(4-fluorophenyl)-4,4-di(trifluoromethyl)cyclopenten-1-yl]benzenesulfonamide;
- 30 5-[2-(4-(aminosulfonyl)phenyl)cyclopenten-1-yl]-1,3-benzodioxole;
- 4-[2-(3,4-dimethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-fluoro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-(3-chloro-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(3-trifluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 5 4-[2-phenylcyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3,4-difluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 10 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 15 4-[2-(3,4-dichlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 20 4-[2-(3-fluoro-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methoxy-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
- 25 4-[2-(3,4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-methyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 30 4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-trifluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 4-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
- 35 4-[2-[3,4-di(trifluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;

- 4-[2-(3-methyl-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-fluoro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
5 4-[2-(3-methoxy-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-chloro-4-trifluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-difluoromethyl-4-methylphenyl)cyclopenten-1-yl]benzenesulfonamide;
10 4-[2-(3-difluoromethyl-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-difluoromethyl-4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;
15 4-[2-(3-difluoromethyl-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-[3,4-di(diifluoromethyl)phenyl]cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methyl-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
20 4-[2-(3-fluoro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide;
4-[2-(3-methoxy-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide; and
25 4-[2-(3-chloro-4-difluoromethylphenyl)cyclopenten-1-yl]benzenesulfonamide.

41. The method of Claim 39 wherein 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
30 or a pharmaceutically-acceptable salt thereof.

42. The method of Claim 39 wherein said compound is 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene, or a pharmaceutically-acceptable
35 salt thereof.

43. The method of Claim 39 wherein said compound is 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide; or a pharmaceutically-acceptable salt thereof.

5

44. The method of Claim 39 wherein said compound is 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide; or a pharmaceutically-acceptable salt thereof.

10

45. The method of Claim 39 wherein said compound is 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide; or a pharmaceutically-acceptable salt thereof.

15

46. The method of Claim 39 wherein said compound is 1-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene, or a pharmaceutically-acceptable salt thereof.

20

47. The method of Claim 37 for use in treatment of inflammation.

48. The method of Claim 37 for use in treatment of an inflammation-associated disorder.

25

49. The method of Claim 48 wherein the inflammation-associated disorder is arthritis.

50. The method of Claim 48 wherein the inflammation-associated disorder is pain.

30

51. The method of Claim 48 wherein the inflammation-associated disorder is fever.

A. CLASSIFICATION OF SUBJECT MATTER

C 07 C 317/14, C 07 C 311/16, C 07 D 213/24,
A 61 K 31/095, A 61 K 31/44

According to International Patent Classification (IPC) or to both national classification and IPC 6

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C 07 C 317/00, C 07 C 311/00, C 07 C 323/00,
C 07 C 405/00, C 07 D 213/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE, A1, 4 212 628 (METZLER et al.) 21 October 1993 (21.10.93), claims 1,6 (cited in the application).	1, 25
A	EP, A2, 0 142 801 (MERCK) 29 May 1985 (29.05.85), abstract.	1, 25, 36
A	PURE AND APPLIED CHEMISTRY, vol. 56, no. 7, issued 1984, W.H. LAARHOVEN, "Aspects of the Photochemistry of Aryl- -Ethylenes", pages 1225-1240, scheme 7	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

06 October 1994

Date of mailing of the international search report

26. 10. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

REIF e.h.

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	(cited in the application). -- INDIAN JOURNAL OF CHEMISTRY, B section, vol. 14B, issued 1976, July, O.P.MALIK et al. "Studies on Cannabinoids: Part III -Synthesis of 9,10,11,11a- -Tetrahydro-6H,8H-pyrido (1,2-c)(1,3)benzoxazine", pages 975-978, especially page 975 (cited in the application). --	1
A	THE JOURNAL OF ORGANIC CHEMISTRY, vol. 36, issued 1971, DAVID Y.CURTIN et al. "Restricted Rotation of Aryl Rings in cis-1,2-Diarylcyclo- pentanes and Diarylmethyl- -cyclobutanes", pages 565-572, especially pages 565,566 (cited in the application). ----	1

ANHANG

zum internationalen Recherchen-
bericht über die internationale
Patentanmeldung Nr.

ANNEX

to the International Search
Report to the International Patent
Application No.

ANNEXE

au rapport de recherche inter-
national relatif à la demande de brevet
international n°

PCT/US 94/08047 SAE 94473

In diesem Anhang sind die Mitglieder
der Patentfamilien der im obenge-
nannten internationalen Recherchenbericht
angeführten Patentedokumente angegeben.
Diese Angaben dienen nur zur Unter-
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family
members relating to the patent documents
cited in the above-mentioned inter-
national search report. The Office is
in no way liable for these particulars
which are given merely for the purpose
of information.

La présente annexe indique les
membres de la famille de brevets
relatifs aux documents de brevets cités
dans le rapport de recherche inter-
national visée ci-dessus. Les renseigne-
ments fournis sont donnés à titre indica-
tif et n'engagent pas la responsabilité
de l'Office.

In Recherchenbericht angeführtes Patentedokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
DE A1 4212628	21-10-93	WD A1 9322262	11-11-93
EP A2 142801	29-05-85	DE C0 3478434	06-07-89
		EP A3 142801	16-07-86
		EP B1 142801	31-05-89
		JP A2 60142932	29-07-85
		US A 4656190	07-04-87